

PRINCIPAL BUNDLES ON PROJECTIVE VARIETIES AND THE DONALDSON-UHLENBECK COMPACTIFICATION

V. BALAJI

ABSTRACT. Let H be a semisimple algebraic group. We prove the semistable reduction theorem for μ -semistable principal H -bundles over a *smooth projective variety* X defined over the field $\mathbb{C}$. When X is a *smooth projective surface* and H is simple, we construct the algebro-geometric Donaldson-Uhlenbeck compactification of the moduli space of μ -semistable principal H -bundles with fixed characteristic classes and describe its points. For large characteristic classes we show that the moduli space of μ -stable principal H -bundles is non-empty.

1. INTRODUCTION

The purpose of this paper can broadly be termed two-fold. Its first objective is to prove the *semistable reduction theorem* for the isomorphism classes of μ -semistable principal bundles (in the sense of Ramanathan-Mumford) with a semisimple structure group H over *smooth projective varieties* X *defined over* $\mathbb{C}$. In fact, we prove the semistable reduction theorem for classes of μ -semistable *quasibundles* (Def 2.1). This generalises in its entirety, the basic theorem of Langton which proves that the functor of isomorphism classes of μ -semistable torsion-free sheaves is *proper*. The approach is a generalisation of the one in [3] and [4], where this theorem is proved for curves.

Carrying out the generalisation to the higher dimensional case involves several new ingredients; for instance, one needs the new notion of *quasibundles* (due to A.Schmitt). This plays the role of the μ -semistable torsion-free sheaf so as to realise the boundary points. The final proof is concluded with key inputs from Bruhat-Tits theory.

Since the proof of the semistable reduction theorem is rather long and complicated, it is probably appropriate at this point to highlight the basic differences between our proof and that of Langton in the case of families of torsion-free sheaves. Except at the very beginning, our proof follows an entirely different path to that of Langton primarily because of the fact that it is not even clear if there is a canonical extension of a family of principal bundles parametrised by a punctured disc, *be it even unstable*, across the puncture. The problem is no longer sheaf theoretic and one is forced to address the problem of torsors with structure group which could be *non-reductive group schemes*. It is to handle this problem that one requires some

aspects of Bruhat-Tits theory. We believe that these new aspects which come up here should also be of general applicability in similar situations where compactification questions need to be addressed for bundles with general structure groups.

More precisely, in his proof Langton first extends the family of semistable torsion free sheaves to a torsion-free sheaf in the limit although non-semistable. In other words, the “structure group” of the limiting bundle over a *big* open subset still remains GL_n . Then by a sequence of *Hecke modifications* he reaches the semistable limit without changing the isomorphism class of the sheaf over the generic fibre. Instead, we are forced to extend the family of semistable *rational* H_K -bundles to a rational H'_A -bundle with the limiting bundle remaining semistable, but the structure group becoming *non-reductive* in the limit (being the closed fibre of the group scheme H'_A). In other words, one loses the reductivity of the structure group scheme. Then by using Bruhat-Tits theory, we relate the group scheme H'_A to the reductive group scheme H_A without changing the isomorphism class of the bundle over the generic fibre as well as the semistability of the limiting bundle.

Let A be a complete discrete valuation ring and let K be its quotient field and $k = \mathbb{C}$ its residue field. Our first main theorem is the following (X is an arbitrary *smooth complex projective variety*):

THEOREM 1.1. (*Theorem 4.1*) *Let $\mathbf{P}_K$ be a family of semistable principal H -quasibundles on $X \times \text{Spec } K$, or equivalently, if H_K denotes the group scheme $H \times \text{Spec } K$, a semi-stable H_K -quasibundle $\mathbf{P}_K$ on X_K . Then there exists a finite extension L/K , with the integral closure B of A in L , such that, $\mathbf{P}_K$, after base change to $\text{Spec } B$, extends to a semistable H_B -quasibundle $\mathbf{P}_B$ on X_B .*

We now turn to the second goal of this paper which is to give an algebro-geometric compactification of the moduli space of μ -semistable principal bundles over *smooth projective surfaces*. In fact, we construct a *reduced projective scheme* which can be termed the Donaldson–Uhlenbeck compactification of the moduli space of μ -stable principal H -bundles for a general simple group H (cf. Corollary 6.18). In the vector bundle case such an algebro-geometric construction was given by J.Li (cf. [24], [25]). (see also J.Morgan ([28])).

The theorem of Ramanathan and Subramaniam ([33]), which is a generalisation Donaldson-Uhlenbeck-Yau theorem to the case of principal bundles, gives an identification of antiselfdual (ASD) Yang-Mills bundles over X with general structure groups with μ -stable principal bundles. Therefore, our construction of the Donaldson-Uhlenbeck compactification can be viewed as a natural compactification of the moduli space of antiselfdual (ASD) Yang-Mills bundles over X with general structure groups. We remark that for the case of principal bundles with ASD connections, even a topological compactification has not been constructed although one can perhaps extract such a construction from the text [12].

More precisely, we have the following theorem:

THEOREM 1.2. (*Theorem 6.19*)

- (1) Let H be a semisimple algebraic group and $\rho : H \hookrightarrow SL(V)$ a faithful rational representation of H . There exists a reduced projective scheme $M_H(\rho)$ which parametrises equivalence classes of μ -semistable H -quasibundles with fixed characteristic classes, on the smooth projective surface X . This has an open subscheme of equivalence classes of μ -semistable principal H -bundles M_H^0 .
- (2) Let H be a simple algebraic group. Then set-theoretically the closure $\overline{M_H^0}$ of μ -semistable principal H -bundles M_H^0 in $M_H(\rho)$ is a subset of the disjoint union:

$$\mathfrak{M}_H = \coprod_{l \geq 0} M_H^{\mu\text{-poly}}(\mathfrak{c} - l) \times S^l(X)$$

where $M_H^{\mu\text{-poly}}(\mathfrak{c} - l)$ is the moduli space of μ -semistable principal H -bundles with characteristic classes $\mathfrak{c} - l$ (represented as classes of polystable bundles); in particular, the big stratum is $M_H^{\mu\text{-poly}}(\mathfrak{c}) = M_H^0$.

- (3) When H is simple the underlying set of points of the moduli space $\overline{M_H^0}$, upto homeomorphism, is independent of the representation ρ .
- (4) Let $m = m_\rho$ be the Dynkin index of the representation ρ (Def 6.1). The canonical morphism $f_\rho : M_H(\rho) \longrightarrow M_{SL(V)}$ maps a copy of $S^l(X) \subset M_h(\rho)$ to the symmetric power $S^{ml}(X) \subset M_{SL(V)}$ by sending any cycle Z to $m \cdot Z$.

In the above theorem, the formal construction of the moduli space is by itself not too difficult. However, the description of its points is quite involved. The method of proof is along the lines of the proof of J.Li (cf. [24]) and the methods in the paper of Le Potier ([23]) (cf. [19] for a lucid treatment of this approach). In the description of the points of the moduli space of H -quasibundles and their relationship with the associated moduli space of $SL(V)$ -bundles, the notion of Dynkin index of the representation ρ makes a natural entry and plays a key role in defining intrinsically the *cycles associated to the points of the boundary of the moduli space*. Its significance has already been noted in the paper by Atiyah, Hitchin and Singer [2] for bundles on the real four-sphere and in the paper by Kumar, Narasimhan and Ramanathan [21] for principal bundles on curves.

Since the construction of the Donaldson–Uhlenbeck compactification even in the case of vector bundles ([24]) is not entirely by the methods of GIT it is only natural that for the general case of arbitrary structure groups which we consider, the use of GIT is only peripheral. We may recall that the methods of GIT give as a consequence the projectivity of the quotient space constructed. From this standpoint, our first theorem (Theorem 4.1) is absolutely essential towards proving the compactness. The approach is to

separate the proof of the properness (by proving the semistable reduction), and the construction of the moduli (by the process of separating points using sections of a suitable determinant line bundle).

In [11], S. Donaldson remarks that it is natural to expect a generalised theory for his polynomial invariants arising from the Yang-Mills moduli for bundles with general structure groups. He also comments that the Uhlenbeck compactness theorem should naturally hold for the case of general structure groups. One could say that this is indeed the case in the light of the semistable reduction theorem mentioned above as well as the description of the points of the moduli which are added to compactify the ASD moduli space. It is further remarked in [11] that if a general theory of these moduli spaces is given, then one expects results such as the *vanishing theorems* for these invariants to hold in this general setting. The existence theorem and the description of points of the moduli in the present paper aims at securing the foundations of a precise theory towards this end.

After this paper was completed, the work of Braverman, Finkelberg and Gaietsgory ([9]) was brought to our attention by Alexander Schmitt. They base themselves on a formulation due to V. Ginzburg. The issues which motivate them in considering Uhlenbeck spaces for principal bundles with arbitrary structure groups are deep and far-reaching but quite distinct from ours. Our paper and [9] both aim at the construction of these spaces, but the methods used are altogether different. The coincidence in terminology of *quasibundles* is also surprising since their notion and ours do not seem to be related. The approach in [9] can broadly be termed *adèlic* and in the setting of curves this has been used earlier in [21].

We believe that in this paper we have in fact settled affirmatively some of the basic questions raised in [9, Page 1], particularly those related to *moduli*. It should be very interesting to establish precise relationships between our paper and [9].

In the final section we prove the following *existence* theorem on which hinges any computation of Donaldson polynomials associated to these moduli spaces.

THEOREM 1.3. (*Theorem 7.9*) *Let H be a semisimple algebraic group over $\mathbb{C}$. Then the moduli space $M_H(c)^s$ of μ -stable principal H -bundles on a smooth projective surface X is non-empty for large c .*

In the case when $H = SL(2)$ this is highly non-trivial and uses some deep ideas; this is due to Taubes [40] and later due to Gieseker [16]. Both methods are deformation theoretic, but the method used by Taubes is differential geometric (*gluing techniques*), whereas Gieseker used degeneration techniques in an algebraic geometric setting to prove the non-emptiness of the moduli space of μ -stable $SL(2)$ -bundles. Our approach for the general case of arbitrary semisimple H is to draw on some classical representation theory, by using what are known as *principal $SL(2)$'s* in a semisimple group. We then construct μ -stable principal H -bundles starting from $SL(2)$ -bundles

for such principal $SL(2)$'s in H . The important point in these existence results is that the bounds are dependent only on $p_g(X)$ and not on the polarisation Θ on X . Note that our theorem implies the non-emptiness of μ -stable $SL(r)$ -bundles for all r .

The proof of this theorem (where we construct H -bundles starting from $SL(2)$ -bundles with $SL(2) \subset H$) and the construction of the moduli space, where we use a faithful representation $H \subset SL(V)$, indicate the strong possibility of an *algebra* of Donaldson polynomials coming from the tensor structure on the category of representations of H .

Very recently, Gómez and Sols ([18]) and Schmitt ([35], [36]) have constructed compactifications of moduli spaces of principal bundles on higher dimensional varieties using the Gieseker–Maruyama approach for torsion-free sheaves. The non-emptiness of these spaces (over surfaces) is also therefore a consequence of our Theorem 7.9.

Gómez and Sols follow and generalise the methods of Ramanathan to higher dimensional varieties. Schmitt gives an alternative approach, via GIT again, for the moduli construction but in either case this means that they work with a “Gieseker–Maruyama” type definition for semistability. Schmitt introduces the concept of *honest singular principal bundles* to recover the boundary points of the moduli space. The *singular bundles* of Schmitt or equivalently our *quasibundles* (Def 2.1) play the key role of giving the boundary points in our moduli space. It seems possible that the moduli spaces that we have constructed can be recovered by a generalised blow-down of the Gómez–Sols moduli but this needs to be investigated.

In contrast, the striking feature that emerges here is that the underlying set of points of our moduli space, (*upto homeomorphism*), is independent of the choice of a representation of the structure group, while the moduli spaces of Gómez–Sols and Schmitt are invariably dependent on the faithful representation chosen. We however make no statement on any natural scheme structure on the moduli space. In fact, this is the case even in the usual Donaldson–Uhlenbeck compactification for vector bundles.

The brief layout of the paper is as follows: Sections 1 to 4 are devoted to the proof of the semistable reduction theorem. Section 5 is devoted to the construction of the compactification and section 6 for the description of its points. Section 7 contains the proof of the non-emptiness of the moduli spaces of stable bundles for large characteristic classes.

Acknowledgments: I am grateful to C.S.Seshadri and D.S.Nagaraj for their patience and help in this work. I thank M.S.Narasimhan, V.Uma and S.Bandhopadhyaya for helpful discussions. I am extremely grateful to Alexander Schmitt for a meticulous reading of an earlier version of this paper. His comments have been very helpful in clarifying many issues in this paper.

CONTENTS

1. Introduction
2. Rational bundles and principal quasibundles

3. Extension of structure group to the flat closure
 4. Semistable reduction for quasibundles over projective varieties
 5. Construction of the moduli space of bundles over surfaces
 6. The geometry of the moduli of H -bundles
 7. Non-emptiness of the moduli space
- References

1.1. Notations and Conventions. Throughout this paper, unless otherwise stated, we have the following notations and assumptions:

- (a) We work over an algebraically closed field k of characteristic zero and without loss of generality we can take k to be the field of complex numbers $\mathbb{C}$.
- (b) X will be a smooth projective variety over k till §4. From §5 onwards it will be a smooth projective surface.
- (c) We fix a hyperplane Θ on X throughout and will use Θ for all degree computations.
- (d) By a **large** or **big** open subset $U \subset X$, we mean a subset such that $\text{codim}_X(X - U) \geq 2$.
- (e) H is a *semisimple* algebraic group, and G , unless otherwise stated will always stand for the general linear group $GL(V)$ for a finite dimensional vector space V . Their representations are finite dimensional and rational.
- (f) A is a discrete valuation ring (which could be assumed to be complete) with residue field k , and quotient field K .
- (g) Let E be a principal G -bundle on $X \times T$ where T is $\text{Spec } A$. If $x \in X$ is a closed point then we shall denote by $E_{x,A}$ or $E_{x,T}$ (resp $E_{x,K}$) the restriction of E to the subscheme $x \times \text{Spec } A$ or $x \times T$ (resp $x \times \text{Spec } K$). Similarly, $p \in T$ will denote the closed point of T and the restriction of E to $X \times p$ will be denoted by E_p .
- (h) In the case of $G = GL(V)$, when we speak of a principal G -bundle we identify it often with the associated vector bundle (and can therefore talk of the degree of the principal G -bundle with respect to the choice of Θ).
- (i) We denote by E_K (resp E_A) the principal bundle E on $X \times \text{Spec } K$ (resp $X \times \text{Spec } A$) when viewed as a principal H_K -bundle (resp H_A -bundle). Here H_K and G_K (resp H_A and G_A) are the product group schemes $H \times \text{Spec } K$ and $G \times \text{Spec } K$ (resp $H \times \text{Spec } A$ and $G \times \text{Spec } A$).
- (j) If H_A is an A -group scheme, then by $H_A(A)$ (resp $H_K(K)$) we mean its A (resp K)-valued points. When $H_A = H \times \text{Spec } A$, then we simply write $H(A)$ for its A -valued points. We denote the closed fibre of the group scheme by H_k .
- (k) Let Y be any G -variety and let E be a G -principal bundle. For example Y could be a G -module. Then we denote by $E(Y)$ the associated bundle with fibre type Y which is the following object: $E(Y) = (E \times Y)/G$ for the twisted action of G on $E \times Y$ given by $g.(e, y) = (e.g, g^{-1}.y)$.
- (l) If we have a group scheme H_A (resp H_K) over $\text{Spec } A$ (resp $\text{Spec } K$) an H_A -module Y_A and a principal H_A -bundle E_A . Then we shall denote the associated bundle with fibre type Y_A by $E_A(Y_A)$.
- (m) By a family of H bundles on X parametrised by T we mean a principal H -bundle on $X \times T$, which we also denote by $\{E_t\}_{t \in T}$.

2. RATIONAL BUNDLES AND PRINCIPAL QUASIBUNDLES

Let X be a smooth projective variety over $\mathbb{C}$. Let $\rho : H \hookrightarrow SL(V)$. Let $\mathcal{E}$ be a torsion-free sheaf on X and let $U(\mathcal{E})$ be the largest open subset where $\mathcal{E}$ is locally free. Let $S_{\mathcal{E}}$ be the affine X -scheme given by $\underline{\text{Spec}} \text{Sym}^*(\mathcal{E} \otimes \mathcal{O}(V))$. There is a canonical action of H on $S_{\mathcal{E}}$ and we consider the categorical quotient $\underline{\text{Spec}} (\text{Sym}^*(\mathcal{E} \otimes \mathcal{O}(V))^H) = (S_{\mathcal{E}})//H$. Since H is assumed to be semisimple, the action has a non-empty collection of semistable points.

Suppose that we are given a morphism (*which we term a reduction section for the obvious reasons*) $\sigma : X \rightarrow (S_{\mathcal{E}})//H$, which on the open subset $U(\mathcal{E})$ gives a *genuine reduction of structure group* to H . We obtain an X -scheme $P \rightarrow X$, by pulling back the quotient map $q : S_{\mathcal{E}} \rightarrow (S_{\mathcal{E}})//H$. This notion is due to Alexander Schmitt, and the pair $(\mathcal{E}, \sigma)$ is termed by him an *honest singular principal bundle*. This is a natural generalisation of the classical notion of a *frame bundle* associated to a vector bundle.

DEFINITION 2.1. (*Rational principal bundles and principal quasibundles*)

- (a) Following ([34]), by a rational principal H -bundle, we mean a principal H -bundle on a large open $U \subset X$.
- (b) (A.Schmitt) By an H -quasibundle, we mean a scheme $P \rightarrow X$ as above. Let $U(P)$ denote the largest open subset of X where P is a genuine principal H -bundle. In particular, $P|_{U(P)}$ is a rational principal H -bundle.

Remark 2.2. We remark that whenever we speak of a quasibundle we have the following collection of objects:

- (a) A faithful representation $H \hookrightarrow SL(V)$.
- (b) A torsion-free sheaf $\mathcal{E}$ with generic fibre of type V^* .
- (c) A diagram:

$$\begin{array}{ccc} P & \longrightarrow & S_{\mathcal{E}} \\ \downarrow & & \downarrow \\ X & \xrightarrow{\sigma} & S_{\mathcal{E}}//H \end{array}$$

Remark 2.3. Since the notion of quasibundles will play a key role in what proceeds, we will briefly recall its salient features, especially those which will be frequently used in this paper.

Let T be an arbitrary normal variety and $\mathcal{E}$ a torsion-free sheaf on T . We can identify the affine T -scheme $S_{\mathcal{E}}$ with the space $\underline{\text{Hom}}(\mathcal{E}, V^* \otimes \mathcal{O}_X)$ and similarly the affine T -scheme $S_{\mathcal{E}}//H$ with the space $\underline{\text{Hom}}(\mathcal{E}, V^* \otimes \mathcal{O}_X)//H$ (cf. [35, 3.7, 3.8]).

Let $\mathcal{U}$ be the maximal open subset of T such that $\mathcal{E}|_{\mathcal{U}}$ is locally free with general fibre V and trivial determinant and let $\rho : H \hookrightarrow GL(V)$. Then one knows that $\mathcal{U}$ is a *big* open subset of T . A reduction of structure group of the principal $GL(V)$ -bundle underlying the vector bundle $\mathcal{E}|_{\mathcal{U}}$ can be viewed as

a section:

$$\sigma_{\mathcal{U}} : \mathcal{U} \longrightarrow \underline{Isom}(\mathcal{E}|_{\mathcal{U}}, V \otimes \mathcal{O}_X)/H$$

Now observe that $\underline{Isom}(\mathcal{E}|_{\mathcal{U}}, V \otimes \mathcal{O}_X)/H \subset S_{\mathcal{E}}//H$ and further, since $S_{\mathcal{E}}//H$ can be embedded as closed subscheme of a vector bundle over T , it follows by Hartogs theorem that the reduction $\sigma_{\mathcal{U}}$ extends uniquely to a section:

$$\sigma : X \longrightarrow S_{\mathcal{E}}//H.$$

Following [32] we have the following definitions.

DEFINITION 2.4. (A. Ramanathan) A rational principal H -bundle E is said to be μ -semistable (resp. μ -stable) if $\forall$ parabolic subgroup P of H , $\forall$ reduction $\sigma_P : X \longrightarrow P(H/P)$ and $\forall$ dominant character χ of P , the bundle $\sigma_P^*(L_{\chi})$ has degree ≤ 0 (resp. < 0). (cf.[32]). We note that in this convention, a dominant character χ of P induces a negative ample line bundle on G/P . Note further that this definition makes sense since the degree of the line bundle is well defined on large open subsets. This definition works for reductive groups as well.

DEFINITION 2.5. A H -quasibundle P is said to be μ -semistable (resp. μ -stable) if the induced rational principal H -bundle $P|_{U(P)}$ is μ -semistable (resp. μ -stable).

DEFINITION 2.6. A reduction of structure group of E to a parabolic subgroup P is called *admissible* if for any character χ on P which is trivial on the center of H , the line bundle associated to the P -bundle E_P obtained by the reduction of structure group, has degree zero.

DEFINITION 2.7. An H -bundle E is said to be *polystable* if it has a reduction of structure group to a Levi subgroup R of a parabolic P such that the R -bundle E_R obtained by the reduction, is stable and the extended P bundle $E_R(P)$ is an admissible reduction of structure group for E . Since the definition involves only degrees of line bundles, it clearly holds good for rational principal bundles as well. (cf. [34])

Remark 2.8. It is clear that we have the natural notions of polystability for H -quasibundles as well. So a quasibundle P is termed polystable if the induced rational principal H -bundle $P|_{U(P)}$ is *polystable*.

2.0.1. Semistability and polystability over curves. The study of semistability and polystability of principal bundles on curves was initiated by A.Ramanathan. Over the years it has developed in many directions and the results which one requires are scattered in the literature. Polystability is also differently called *quasi-stability* in the literature but we avoid this terminology for the obvious reasons (we have already a notion of quasibundles ..).

For the convenience of the reader we gather some of the relevant facts with appropriate references in the following theorem:

THEOREM 2.9. *Let C be a smooth projective curve over the field of characteristic zero. Let H a semisimple algebraic group.*

The following are equivalent:

- (i) *A principal H -bundle E is polystable in the sense of Ramanathan (Def 2.7).*
- (ii) *There exists a faithful representation $H \hookrightarrow GL(V)$ such that the associated bundle $E(V)$ is polystable of degree 0.*
- (iii) *For every representation $H \rightarrow GL(W)$, the bundle $E(W)$ is polystable of degree 0.*
- (iv) *Let $ad : H \rightarrow GL(\mathfrak{h})$ be the adjoint representation. Then $E(\mathfrak{h})$ is polystable (of degree zero).*
- (v) *The bundle E arises from a representation $\rho : \pi_1(C) \rightarrow K$, where K is a maximal compact subgroup of H .*
- (vi) *The bundle E carries an Einstein–Hermitian connection.*

Proof: The equivalence of (i) and (v) is the main theorem of Ramanathan, which generalises the Narasimhan–Seshadri theorem for principal bundles. ([31]).

(i) $\Leftrightarrow$ (vi) is the main theorem of [33]. The equivalence (iv) $\Leftrightarrow$ (v) is shown in [1, Lemma 10.12].

By [33, Prop 1], we may go modulo the center and assume that the group H has trivial center. Therefore, the adjoint representation $ad : H \hookrightarrow GL(\mathfrak{h})$ is a faithful representation. From this standpoint, (ii), (iii) and (iv) are equivalent by a Tannakian principle and the proof can be found in [3, Prop 2.3]. The argument there is for semistability, but the changes needed to be made for polystability are easy since all bundles involved are of degree 0 (see also the proof of Prop 3.11 below).

q.e.d

Remark 2.10. In fact, the equivalence [iv] $\Leftrightarrow$ [v] holds more generally even for H reductive (cf. [1, Lemma 10.12]).

2.1. Some key lemmas. We recall the following couple of facts about torsion-free sheaves which we will use in this work. The first one is rather well known.

Let $C \subset U$ be a smooth projective curve. We recall:

LEMMA 2.11. (cf. [3, Lemma 2.10]) *Let $T = \text{Spec } A$ and let E_T be a family of vector bundles on $C \times T$ such that E_p is semistable of degree 0. Let s_K be a section of the family E_K restricted to $C \times \text{Spec } K$, with the property that for a base point $x \in C$, the section s_K extends along $x \times T$ to give a section of $E_{x \times T}$. Then the section extends to the whole of $C \times T$.*

We have more generally:

LEMMA 2.12. *Let W be a family of semistable vector bundles with $c_1 = 0$ on U parametrised by T , i.e a vector bundle on U_A , where $U \subset X$ is a large open*

subset of X . Suppose that we are given a regular section $s_K : U_K \longrightarrow W_K$ such that, for an irreducible smooth divisor $Y \subset U$, the section extends as a regular section along Y_A . Then the section s_K extends to a regular section s_A on U_A .

Proof: For the section s_{T-p} , viewed as a section of W_{T-p} we have two possibilities, since U_A is normal and the polar set is a divisor:

- (a) it either extends as a regular section s_T .
- (b) or it has a pole along $U \times p$.

By the given property, we have a section $\sigma_Y : Y \times T \longrightarrow W_Y$ with the property that, at $\forall x \in Y$, $s_t(x) = \sigma_Y(x, t) \forall t \in T - p$;

So to complete the proof, we need to check that the possibility (b) cannot hold:

Suppose it does hold. Then the section $s_{T-p} = s_K$ is a section of W_K , i.e, a *rational section* of W with a pole along the divisor $U \times p \subset U \times T$, of order $k \geq 1$.

Thus by multiplying s_{T-p} by π^k we get a regular section s'_T of W on $U \times T$. If $s'_T = \{s'_t\}_{t \in T}$, then we have:

- (i) $s'_t = \lambda(t) \cdot s_t$, $t \in T - p$ where $\lambda : T \longrightarrow \mathbb{C}$ is a function given by π^k , having zeros of order k at p .
- (ii) s'_p is a non-zero section of W_p . Here we first note that W_p is a bundle on U which is large and hence the μ -semistability is completely determined by W_p . Further, by taking W_p^{**} , this extends as a reflexive sheaf to the whole variety X and remains μ -semistable of degree 0. By Hartogs' theorem (since reflexive sheaves are *normal*), the section s'_p extends to a non-zero section of W_p^{**} to the whole of X .

Since $\mathcal{O}_X$ is stable and degree 0, it follows that s'_p gives a short exact sequence of sheaves on X :

$$0 \longrightarrow \mathcal{O}_X \longrightarrow W_p^{**} \longrightarrow Q \longrightarrow 0$$

where the torsion part of Q is supported in a subset of codimension ≥ 2 (if the quotient sheaf had a divisor in its support then by taking determinants, we see that there is a contradiction to the equality of the degrees $\deg(\mathcal{O}_X) = \deg(W_p^{**}) = 0$).

Thus it follows that there exists

$$y \in Y \text{ such that } s'_p(y) \neq 0. \quad (*)$$

By the assumption that the section extends along Y_A , and by the fact that $y \in Y$ it follows that $s_t(y) = \sigma_Y(y, t)$, for $t \in T - p$, and hence

$$s'_t(y) = \lambda(t) \cdot \sigma_Y(y, t) \text{ for } t \in T - p.$$

Therefore, by continuity, since $\sigma_Y(y, p)$ is well-defined, we see that $\lambda(t) \cdot \sigma_Y(y, t)$ tends to $\lambda(p) \cdot \sigma_Y(y, p) = 0$, as $t \longrightarrow p$.

Also, $s'_t(y) \longrightarrow s'_p(y)$ since $t \longrightarrow p$. Hence by continuity again, it follows that $s'_p(y) = 0$, which contradicts (*).

Thus the possibility (b) does not occur and we are done.

q.e.d

2.1.1. *A Refinement of Langton's theorem.* Let $R^{\mu ss}$ denote the subset of the usual Quot scheme which parametrises the usual torsion-free sheaves with fixed topological type and Hilbert polynomial. Let $\mathcal{G}$ be the group $SL(\chi)$ (χ being the Euler characteristics of sheaves in $R^{\mu ss}$) acting on the (open subset) $R^{\mu ss}$ of the Quot scheme.

LEMMA 2.13. (*Modified Langton*) Let E_K denote a family of μ -semistable torsion-free sheaves of degree zero on $X \times \text{Spec } K$. Then, (by going to a finite cover S of T if need be) the sheaf E_K extends (upto isomorphism) to a family E_A with the property that the limiting sheaf E_0 is in fact polystable.

Proof. By definition E_K gives a K -valued point $x_K : \text{Spec}(K) \longrightarrow R^{\mu ss}$

By Langton, there exists an A -valued point $x_A : \text{Spec}(A) \longrightarrow R^{\mu ss}$ such that x_p is given by E_p , which is a semistable torsion-free sheaf.

One knows that (for a choice of a Jordan–Holder filtration) there exists a family $F_t|_{t \in \mathbb{A}}$ such that $F_t \simeq E_p$ for $t \neq 0$ and the limit is a polystable sheaf $F_0 \simeq gr^\mu(E_p)$. Let us denote this family by a morphism $f : \mathbb{A} \longrightarrow R^{\mu ss}$ and the point $f(0) = y_0$ for $t = 0$.

By going to a finite cover if need be, we may assume that there exists a point $g \in \mathcal{G}(\mathbb{C}(t))$ such that $g \cdot E_p \simeq F_{\mathbb{C}(t)}$ with limit given by F_0 .

Let $D = \overline{\mathcal{G} \cdot x_K}$ (the $\mathcal{G}(K)$ -orbit closure in $R^{\mu ss}$) considered as a $\mathbb{C}$ -variety. Then by definition the point $x_p \in D$ (where x_p corresponds to the sheaf E_p). Further, D is $\mathcal{G}$ -stable. Therefore, $g \cdot x_p \in D(\mathbb{C}(t))$. Since D is closed it follows that the point y_0 belongs to D .

Again, since D is irreducible, we can join y_0 to the orbit $\mathcal{G} \cdot x_K$ and we get a scheme S , the spectrum of a discrete valuation ring, such that $s_0 = y_0$ and $S - s_0 = \text{Spec}(L)$ where L is a finite extension of K (consider the natural map $q : \mathcal{G} \cdot x_K \longrightarrow \text{Spec}(K)$. Therefore $(S \cap \mathcal{G} \cdot x_K)$ is mapped to $\text{Spec}(K)$ by q which gives the extension L/K .)

We also get a resulting family $\{E'_s\}_{s \in S}$ such that $E'_s \simeq E_{q(s)}$ for $s \neq 0$ and $E'_{s_0} \simeq gr^\mu(E_p)$. We are done.

q.e.d

3. EXTENSION OF STRUCTURE GROUP TO THE FLAT CLOSURE

Fix a faithful representation $\rho : H \hookrightarrow G$ defined over $\mathbb{C}$. Consider the extension of structure group of the principal H_K -quasibundle (E_K, σ_K) via the induced K -inclusion $\rho_K : H_K \hookrightarrow G_K$. In other words, on a large open U_K , we are given a reduction of structure group of the principal G_K -bundle $E_K|_{U_K}$ to H_K .

Then, since $G = GL(n)$, by the properness of the functor of semistable torsion-free sheaves (the Theorem of Langton), there exists a *semistable extension* of E_K to a torsion-free sheaf on $X \times \text{Spec } A$, which we denote by E_A . Call the restriction of E_A to $X \times p$ (identified with X) the *limiting bundle* of E_A and denote it by E_p (as in §1). One has in fact slightly more, which is what we need.

3.1. *Mehta–Ramanathan restriction theorems.* Fix an integer $c \in [1, d-1]$ where $d = \dim(X)$ and

$$(a_1, \dots, a_c) \in \mathbb{Z}^c$$

with $a_i \geq 2$ for all $i \in [1, c]$. For any integer $m \geq 1$, by a complete intersection of type (m) we mean a complete intersection of divisors:

$$D_1 \cap D_2 \cap \dots \cap D_c$$

with $D_i \in |a_i^m \Theta|$.

By a general complete intersection subvariety of type (m) we mean the complete intersection formed by the D'_j s from a non-empty open subset S of the linear system $\prod_{i=1}^{n-1} |a_i^m \Theta|$.

Note that for $m \gg 0$, a *general complete intersection of type (m)* is a smooth irreducible subvariety of X of dimension $d - c$.

We recall the main theorem of [27]:

THEOREM 3.1. (Mehta–Ramanathan). *Let V be a semi-stable (resp. stable, polystable) rational vector bundle over $U \subset X$. Then for all $m \gg 0$ the restriction $V|_{Y^\circ}$ to $Y \cap U$, for a general complete intersection Y of type (m) is a semi-stable (resp. stable, polystable) rational vector bundle. Conversely, if $V|_C$ to a general complete intersection curve of type (m) is semi-stable (resp. stable, polystable) then so is V .*

Remark 3.2. Consider the extended sheaf E_A obtained above with the property that E_p is a polystable torsion-free sheaf of degree 0. Since E_p is a *polystable* torsion-free sheaf of degree 0, let $U = U(E_p) \subset X$ be the largest open set where it is locally free. By the Restriction theorem of Mehta–Ramanathan, for a large m , there is an open subset $S \subset \prod_{i=1}^{n-1} |a_i^m \Theta|$, such that if $Y \in S$ is a complete intersection subvariety, then $E_p|_{Y \cap U}$ remains polystable of degree 0.

Let the $'$ denote intersection with U . We can therefore choose a chain of subvarieties:

$$C \subset \dots (D_1 \cap D_2 \cap \dots \cap D_c)' \dots \subset D_1' \subset U \quad (3.2)$$

and the point $x \in C$ such that the restriction $E_p|_{(D_1 \cap D_2 \cap \dots \cap D_c)'}'$ for every c is *locally free and remains polystable* of degree 0.

Since A is a discrete valuation ring, if we choose an open subset $U \subset X$ where the limiting sheaf E_p is locally free, then the family E_A is also *locally free* when restricted to the large open subscheme U_A (this is easy to see. cf. for example Lemma 5.4 [30]). Hence, if we further restrict E_A to the

curve $C_A \subset U_A$, we get a family of locally free sheaves $E_A|_{C_A}$ on the smooth projective curve C parametrised by $\text{Spec} A$. Further, by choice, the limiting bundle E_p is polystable on C of degree 0.

3.2. *The Flat closure.* We observe the following:

- Note that giving a reduction of structure group of the G_K -bundle E_K on a large U_K is equivalent to giving a section s_K of $E_K(G_K/H_K)$ over U_K .
- Let

$$C \subset \cdots (D_1 \cap D_2 \cap \cdots \cap D_c)' \cdots \subset D_1' \subset U = U(E_p)$$

be as in (3.2) above. We fix a base point $x \in C \subset U$ and denote by $x_A = x \times \text{Spec} A$, the induced section of the family (which we call the *base section*):

$$X_A \longrightarrow \text{Spec} A$$

Since $x \in U$, it allows us to work with genuine principal bundles and their restrictions to the section x_A defined by the base point x .

- From now on, unless otherwise stated, we shall fix this chain of smooth subvarieties of U along with the base point x .
- Let $E_{x,A}$ (resp $E_{x,K}$) be as in §1, the restriction of E_A to x_A (resp x_K). Thus $s_K(x)$ is a section of $E_K(G_K/H_K)_x$ which we denote by $E_x(G_K/H_K)$.
- Since $E_{x,A}$ is a principal G -bundle on $\text{Spec} A$ and therefore trivial, it can be identified with the group scheme G_A itself. *For the rest of the article we fix one such identification, namely:*

$$\xi_A : E_{x,A} \longrightarrow G_A.$$

- Since we have fixed ξ_A we have a canonical identification

$$E_x(G_K/H_K) \simeq G_K/H_K$$

which therefore carries a natural *identity section* e_K (i.e the coset $id.H_K$). Using this identification we can view $s_K(x)$ as an element in the homogeneous space G_K/H_K .

- Let $\theta_K \in G(K)$ be such that $\theta_K^{-1} \cdot s_K(x) = e_K$. Then we observe that, the isotropy subgroup scheme in G_K of the section $s_K(x)$ is $\theta_K.H_K.\theta_K^{-1}$. (We remark that such a θ_K will exist after going to a finite extension of K . By an abuse of notation we will continue calling this extension as K . This is required since $G_K \longrightarrow G_K/H_K$ need not be locally trivial).
- On the other hand one can realise $s_K(x)$ as the identity coset of $\theta_K.H_K.\theta_K^{-1}$ by using the following identification:

$$G_K/\theta_K.H_K.\theta_K^{-1} \xrightarrow{\sim} G_K/H_K.$$

$$g_K(\theta_K.H_K.\theta_K^{-1}) \longmapsto g_K\theta_K.H_K.$$

DEFINITION 3.3. Let H'_K be the subgroup scheme of G_K defined as:

$$H'_K := \theta_K \cdot H_K \cdot \theta_K^{-1}.$$

Using ξ_A we can have a canonical identification:

$$E_x(G_K/H'_K) \simeq G_K/H'_K.$$

Then we observe that, using the above identification we get a section s'_K of $E_K(G_K/H'_K)$, with the property that, $s'_K(x)$ is the *identity section* and moreover, since we have conjugated by an element $\theta_K \in G_A(K)(= G(K))$, the isomorphism class of the H_K -bundle P_K given by s_K does not change by going to s'_K .

Thus in conclusion, the G_A -bundle E_A has a reduction to H'_K given by a section s'_K of $E_K(G_K/H'_K)$, with the property that, at the given base section $x_A = x \times \text{Spec } A$, we have an equality $s'_K(x_A) = e'_K$, with the *identity element* of G_K/H'_K (namely the coset $id \cdot H'_K$).

DEFINITION 3.4. The *flat closure* of the reduced group scheme H'_K in G_A is defined to be the schematic closure of H'_K in G_A with the reduced scheme structure. Let H'_A denote the flat-closure of H'_K in G_A .

Remark 3.5. Let I_K be the ideal defining the subgroup scheme H'_K in $K(G)$ (note that G_A (resp G_K) is an affine group scheme and we denote by $A(G)$ (resp $K(G)$) its coordinate ring). If we set $I_A = I_K \cap A(G)$, then it is easy to see that since we are over a discrete valuation ring, I_A is in fact the ideal in $A(G)$ defining the flat closure H'_A .

We then have a canonical identification via ξ_A :

$$E_x(G_A/H'_A) \simeq G_A/H'_A.$$

One can easily check that H'_A is indeed a subgroup scheme of G_A since it contains the identity section of G_A , and moreover, it is faithfully flat over A . Notice however that H'_A need not be a *reductive* group scheme; that is, the special fibre H_k over the closed point need not be reductive.

Observe further that $s'_K(x)$ extends in a trivial fashion to a section $s'_A(x)$, namely the *identity coset section* e'_A of $E_x(G_A/H'_A)$ identified with G_A/H'_A .

3.3. *Chevalley embedding of G_A/H'_A .* As we have noted, H'_A need not be reductive and the rest of the proof is to get around this difficulty. Our first aim is to prove that the structure group of the bundle $E_A(G_A)$ can be reduced to H'_A which is the statement of Proposition 3.7.

We need the following generalisation of a well-known result of Chevalley from [3].

LEMMA 3.6. *There exists a finite dimensional G_A -module W_A such that $G_A/H'_A \hookrightarrow W_A$ is a G_A -immersion.*

3.4. *Extension to flat closure and local constancy.* Recall that the section $s'_K(x)$ extends along the base section x_A , to give $s'_A(x) = w_A$. The aim of this section is to prove the following key theorem.

THEOREM 3.7. *There exists a large open subset $U^o \subset U$ such that the section s'_K extends to a section s'_A of $E_A(G_A/H'_A)$ when we restrict E_A to the open subset $U_A^o \subset U_A$. In other words, the structure group of $E_A|_{U_A^o}$ can be reduced to H'_A ; in particular, if H'_k denotes the closed fibre of H'_A , then the structure group of E_k can be reduced to H'_k .*

Towards this we need the following key result.

PROPOSITION 3.8. *Let E be a polystable principal G -bundle of degree zero on a smooth projective curve C (here $G = SL(n)$ or $GL(n)$). Let W be a G -module and N a G -subscheme of W of the form G/H' .*

If s is a section of $E(W)$ such that for some $x \in C$, $s(x)$ lies in the fibre $E(N)_x$ of the fibration $E(N)$ at $x \in C$, then the entire image of s lies in $E(N)$. Consequently, E has a reduction of structure group to the subgroup H' .

Proof. The bundle E being polystable, it is defined by a “unitary” representation

$$\chi : \pi_1(C) \longrightarrow G$$

which maps into a maximal compact subgroup of G . This implies that if the universal covering $j : Z \longrightarrow C$ is considered as a principal fibre space with structure group $\pi_1(C)$, then the principal G -bundle E is the associated bundle through χ .

Let $\rho : G \longrightarrow GL(W)$ be the representation defining the G -module W . Then $E(W)$ can be considered as the bundle associated to the principal bundle $j : Z \longrightarrow C$ through the representation

$$\rho \circ \chi : \pi_1(C) \longrightarrow GL(W).$$

which maps into the unitary subgroup of $GL(W)$.

By generalities on principal bundles and associated constructions, since

$$E(W) \simeq Z \times^{\pi_1(C)} W$$

a section of $E(W)$ can be viewed as a $\pi_1(C)$ -map

$$s_1 : Z \longrightarrow W$$

Now, since Z is the universal cover of the curve C and s is a section of $E(W)$, then one knows (cf. [29]) that there exists a $\pi_1(C)$ -invariant element $w \in W$ such that s is defined by a map

$$s_1 : Z \longrightarrow W$$

given by $s_1(x) = w$, $\forall x \in C$, i.e “the constant map sending everything to w ”.

Since $w \in W$ is a $\pi_1(C)$ -invariant vector and the action of $\pi_1(C)$ is via the representation χ , we see that χ factors via

$$\chi_1 : \pi_1(C) \longrightarrow H'$$

since $H' = \text{Stab}_G(w)$.

In particular, we get the H' -bundle from the representation χ_1 and clearly this H' -bundle is the reduction of structure group of the G -bundle E given by the section s .

By the very construction of the reduction, the induced H' -bundle is *flat* and also semistable since it comes as the reduction of structure group of the polystable bundle E (by Def 3.9 below). This proves the Proposition 3.8.

q.e.d

DEFINITION 3.9. Let H' be an affine algebraic group not necessarily reductive. Let P be a principal H' -bundle on C a smooth projective curve. We define P to be *polystable* if:

- (a) It is *flat* (in the sense that it comes from the representation χ of the fundamental group of C)
- (b) there exists a faithful representation

$$\rho : H' \longrightarrow GL(V)$$

such that the associated vector bundle $P(V)$ is **polystable of degree zero**).

Remark 3.10. This definition is ad hoc and made only to suit our needs. Let $\mathcal{M} := \mathcal{M}(\rho \circ \chi)$ denote the Zariski closure of the image of $\rho \circ \chi : \pi_1(C) \longrightarrow GL(V)$ in $GL(V)$. We term this the “monodromy” subgroup associated to the representation $\rho \circ \chi$.

Since the bundle associated to $\rho \circ \chi : \pi_1(C) \longrightarrow GL(V)$ i.e $P(V)$, is assumed to be polystable of degree 0, by the Narasimhan–Seshadri theorem, the representation $\rho \circ \chi$ is unitary and also the monodromy subgroup $\mathcal{M}$ is *reductive* (possibly non-connected). This can be viewed as a Tannakian interpretation of polystability. Further, since χ maps into H' , the monodromy subgroup $\mathcal{M}$ is a subgroup of H' .

Let us denote the inclusion of the monodromy subgroup by:

$$\iota : \mathcal{M} \hookrightarrow H'$$

The fact that the bundle P comes from an associated construction via the homomorphism χ implies that P has a reduction of structure group to $\mathcal{M}$. Let the resulting $\mathcal{M}$ bundle be denoted by $P_{\mathcal{M}}$

PROPOSITION 3.11. *Let H' be an affine algebraic group (not necessarily reductive), as above, and let P be a polystable principal H' -bundle on a smooth projective curve C . Let $f : H' \longrightarrow H$ be a morphism from H' to a semisimple group H . Then the associated principal H -bundle $P(H)$ is also polystable.*

Proof. Since P has a reduction of structure group to $\mathcal{M}$, we may view the principal H bundle $P(H)$ as obtained from the homomorphism $f \circ \iota : \mathcal{M} \rightarrow H$. Thus $P(H)$ can be identified with $P_{\mathcal{M}}(H)$.

To check the polystability of the principal H -bundle $P(H)$ by Theorem 2.9, we need only check that if $\psi : H \rightarrow GL(W)$ is any representation of H then the associated bundle $P(H)(W)$ is polystable. In other words we need to check that if $\gamma : \mathcal{M} \rightarrow GL(W)$ (with $\gamma = \psi \circ f \circ \iota$) then the associated bundle $P_{\mathcal{M}}(W) = P(H)(W)$ is polystable.

Observe that by Def 3.9 we have a faithful representation $GL(V)$ of $\mathcal{M}$ (from that of H') such that $P_{\mathcal{M}}(V)$ is polystable of degree zero. Further, we are over a field of characteristic zero and so the $\mathcal{M}$ -module W can be realised as a direct summand of a direct sum of some $T^{a,b}(V) = V^{\otimes a} \otimes V^{*\otimes b}$ (cf. for e.g [39] pp 86. We need the reductivity of $\mathcal{M}$ here, otherwise in general this is only a subquotient and not a direct summand).

Hence the vector bundle $P_{\mathcal{M}}(W)$ is a direct summand of $\bigoplus T^{a,b}(P_{\mathcal{M}}(V))$. Since $P_{\mathcal{M}}(V)$ is polystable of degree zero, so is $T^{a,b}(P_{\mathcal{M}}(V))$.

By assumption, since $P_{\mathcal{M}}$ is *flat* the associated vector bundle via any representation is of *degree zero*. Hence $P_{\mathcal{M}}(W)$ has degree 0.

Since all bundles have degree zero and $\bigoplus T^{a,b}(P_{\mathcal{M}}(V))$ is a direct sum of stable bundles of degree 0 it is easy to check that $P_{\mathcal{M}}(W)$ is also direct sum of stable bundles of degree 0 and hence polystable.

q.e.d

Remark 3.12. The polystability of $P_{\mathcal{M}}(W)$ also follows directly from that of $P_{\mathcal{M}}$ by Remark 2.10. We cannot use the Theorem 2.9 as it stands since $\mathcal{M}$ is in general only reductive. But the advantage here is that the bundle $P_{\mathcal{M}}$ comes from a representation of the fundamental group.

After this brief interlude on curves, we now return to the general setting of higher dimensional varieties X .

PROPOSITION 3.13. *Assume that $\dim(X) = 2$. Let E be a rational polystable principal G -bundle of degree zero on $U \subset X$. Let W be a G -module and N a G -subscheme of W of the form G/H' where $H' = \text{Stab}_G(w)$ for some $w \in W$. If s is a section of $E(W)$ such that for $x \in C \subset U$, the image $s(x) = w$ in the fibre of $E(N)$ at $x \in X$. Then there exists a big open subset $U^o \subset U$, such that the entire image of U^o under s lies in $E(N)$. (Here $C \subset U$ is as in (3.2) where the base point x was chosen in a general C). In particular, E has a reduction of structure group to the subgroup H' on a big open subset.*

Proof: By choice, since $x \in C \subset U$, and $E|_C$ remains polystable of degree 0, it follows by Prop 3.8 that $s(C) \subset N$.

Now, again by the Mehta-Ramanathan theorem for $m \gg 0$, there exists an open $\Omega_m \subset |m\Theta|$ such that for $t \in \Omega_m$, the restriction of $E|_{Y_t}$ to the smooth projective curve Y_t remains *polystable* of degree 0. Further, the set

$\{t \in \Omega_m \mid Y_t \cap C \neq \emptyset\}$ is an open subset of Ω_m , which we continue to call Ω_m . Since $Y_t \cap C \neq \emptyset$, for each $t \in \Omega_m$, there is a point $x_t \in Y_t \cap C$ and hence $s(x_t) \in Y_t$. Thus applying Prop 3.8 to the restriction $E|_{Y_t}$, we see that $s(Y_t) \subset N$ for all $t \in \Omega_m$. Hence, the dense subset $U_1 = \bigcup Y_t \subset U$ is mapped into N i.e $s(U_1) \subset N$. This implies that the entire image $s(U)$ lies in the closure of the orbit $\overline{N} \subset W$.

Now observe that $N \subset \overline{N}$ is an open subset, therefore, it follows that the set of points $U^o = \{u \in U \mid s(u) \in N\}$ is an open subset of U .

We claim that $U^o \subset U$ is also *large*. For if not, then the complement of U_2 in U contains a curve and there are $t \in \Omega_m$ such that this curve in the complement meets Y_t , a contradiction since the entire curve Y_t gets mapped into N . This proves our claim.

q.e.d

Now let X be an smooth projective variety of dimension d .

Since E is a rational polystable bundle there is a *big* U where it is a principal H -bundle. We work with the chosen complete intersection as in (3.2) in such a manner that $C \subset U$. With these choices and the point $x \in C \subset U$ we have the following:

PROPOSITION 3.14. *Let E be a rational polystable principal G -bundle of degree zero on $U \subset X$. Let W be a G -module and N a G -subscheme of W of the form G/H' where $H' = \text{Stab}_G(w)$ for some $w \in W$. Let s is a section of $E(W)$ (defined on U) such that the image $s(x)$, of the point $x \in C \subset U$, lies in the fibre of $E(N)$ at $x \in X$. Then there exists a big open subset $U^o \subset U$, such that the entire image of U^o under s lies in $E(N)$. In particular, E has a reduction of structure group to the subgroup H' on a big open subset.*

Proof: By choice, since $x \in C \subset U$, and $E|_C$ remains polystable of degree 0, it follows by Prop 3.8 that $s(C) \subset E(N)$.

Let us, for the present, denote the divisor D_1 simply by D . By an induction on the dimension d , and Prop 3.13, we see that since $C \subset D' = D \cap U$, and since $E|_{D'}$ is also polystable, there exists a big open subset $D^o \subset D'$ such that $s(D^o) \subset E(N)$.

Let $Y_j \in S$ be any other general divisor. Then, by the property of S , we see that $D \cap Y_j$ is a smooth divisor of D and Y_j . Further, by other choices of $Y_i \in S$ we see that there exists a smooth projective curve $C_j \subset Y_j \cap D$.

Since $D^o \subset D'$ is a *big* open subset, in fact, it is easy to see that, by going to a smaller open subset of S if need be (and fixing it), we can make sure that $C_j \subset D^o$ for all $Y_j \in S$ (cf. 4.4, [27]).

Again $C_j \subset (Y_j \cap U)$ and $E|_{(Y_j \cap U)}$ is also a polystable rational bundle. Since $s(C_j) \subset s(D^o) \subset E(N)$, it follows again by induction on dimension that there exists a big open $Y_j^o \subset Y_j$ such that $s(Y_j^o) \subset E(N)$ for all $Y_j \in S$.

Now as in Prop 3.13, we see that the set of points $U^o = \{u \in U | s(u) \in E(N)_u\}$ is a non-empty open subset of U .

We claim that $U^o \subset U$ is also *large*. For if not, then the complement of U^o in U contains a divisor $R \subset U$ and there are $Y_t \in S$ such that $R \cap Y_t \neq \emptyset$. Moreover, by Bertini's theorem one has $\dim(R \cap Y_t) = \dim R - 1 = \dim(Y_t) - 1$. This implies that $(R \cap Y_t) \not\subset (Y_t - Y_t^o)$, i.e $R \cap Y_t^o \neq \emptyset$. That is, there exists $y \in R \cap Y_t^o$ such that $s(y) \in E(N)_y$ which contradicts the fact that $R \subset U - U^o$.

q.e.d

Remark 3.15. We work with this U^o from now on and since there is no confusion we will call this U .

3.5. Completion of proof of Theorem 3.7. We work over the large open U_A and all bundles in this proof are over U_A

By Lemma 3.6, we have an immersion

$$E_A(G_A/H'_A) \subset E_A(W_A).$$

The given section s'_K of $E_K(G_K/H'_K)$ therefore gives a section u_K of $E(W_K)$.

Recall also that we have a chosen complete intersection

$$C \subset \cdots (D_1 \cap D_2 \cap \cdots \cap D_c)' \cdots \subset D'_1 \subset U$$

and the point $x \in C$ (cf. Remark 3.2).

Further, $u_K(x)$, the restriction of u_K to $x \times T^*$, extends to give a section $u_A(x)$ of $E_x(W_A)$ (restriction of $E_A(W_A)|_{C_A}$ to $x \times T$).

Thus by Lemma 2.11, and by the semistability of $E(W_A)_p$, on C the section u_K extends to give a section u_A of $E(W_A)|_{C_A}$ over $C \times T$.

Again, by Lemma 2.12, we see, by an induction on dimension, that the section u_K which is defined on U_K extends to section u_A of $E_A(W_A)$ over the entire open subscheme U_A .

Now, to prove the Theorem 3.7, we need to make sure that:

The image of this extended section u_A actually lands in $E_A(G_A/H'_A)$.
(*)

This would then define s'_A .

To prove (*), it suffices to show that $u_A(U \times p)$ lies in $E_A(G_A/H'_A)_p$ (the restriction of $E_A(G_A/H'_A)$ to $U \times p$).

Observe that, $u_A(x \times p)$ lies in $E_A(G_A/H'_A)_p$ since $u_A(x) = s'_A(x) = w_A$.

Observe further that, if E_p denotes the principal G -bundle on U , which is the restriction of the G_A -bundle E_A on $U \times T$ to $U \times p$, then $E_A(W_A)_p = E_A(W_A)|(U \times p)$, and we also have

$$\begin{array}{ccc}
E_A(G_A/H'_A)_p & \xrightarrow{\cong} & E_p(G_k/H'_k) \\
\downarrow & & \downarrow \\
E_A(W_A)_p & \xrightarrow{\cong} & E_p(W)
\end{array}$$

and the vertical maps are inclusions:

$$E_A(G_A/H'_A)_p \hookrightarrow E_A(W_A)_p, \text{ and } E_p(G_k/H'_k) \hookrightarrow E_p(W)$$

where $E_p(W) = E_p \times^{H'_k} W$ with fibre as the G -module $W = W_A \otimes k$. Note that G/H'_k is a G -subscheme Y of W .

Recall that E_p is polystable of degree zero. Then, from the foregoing discussion, the assertion that $u_A(U \times p)$ lies in $E_A(G_A/H'_A)$, is a consequence of Proposition 3.14 applied to E_p (we might have to throw away a subset of U of codimension ≥ 2 for this).

Thus we get a section s'_A of $E_A(G_A/H'_A)$ on $U \times T$, which extends the section s'_K of $E_A(G_A/H'_A)$ on $U \times T^*$. This gives a reduction of structure group of the G_A -bundle E_A on $U \times T$ to the subgroup scheme H'_A and this extends the given bundle E_K to the subgroup scheme H'_A .

q.e.d

4. SEMISTABLE REDUCTION FOR QUASIBUNDLES OVER PROJECTIVE VARIETIES

The aim of this section is to prove the following theorem.

THEOREM 4.1. *Let P_K be a family of semistable principal H -quasibundles on $X \times \text{Spec } K$, or equivalently, if H_K denotes the group scheme $H \times \text{Spec } K$, a semi-stable H_K -quasibundle P_K on X_K . Then there exists a finite extension L/K , with the integral closure B of A in L , such that, P_K , after base change to $\text{Spec } B$, extends to a semistable H_B -quasibundle P_B on X_B .*

4.1. Potential good reduction. We begin by observing that we have extended the original rational H_K -bundle upto isomorphism to a rational H'_A -bundle. To complete the proof of the Theorem 4.1, we need to extend the H_K -bundle to an H_A -bundle.

Remark 4.2. We note that in general the group scheme H'_A obtained above need not be a smooth group scheme over A . But in our case since the characteristic of the base field is zero and since H'_A is flat, it is also smooth over A .

Recall that H_A denotes the reductive group scheme $H \times \text{Spec } A$ over A . We need the following crucial result from [3]. It involves key inputs from Bruhat–Tits theory. For details see Appendix of [4].

PROPOSITION 4.3. *There exists a finite extension L/K with the following property: If B is the integral closure of A in L , and if H'_B are the pull-back group schemes, then we have a morphism of B -group schemes*

$$H'_B \longrightarrow H_B$$

which extends the isomorphism $H'_L \cong H_L$.

We also need the following:

LEMMA 4.4. *Let H' be a non-reductive group and let $\psi : H' \longrightarrow H$ be a homomorphism of algebraic groups. Let P be a rational principal H' -bundle on the big open subset $U \subset X$ obtained as reduction of structure group of a polystable vector bundle E of degree zero through an inclusion $H' \hookrightarrow GL(V)$. Then, by the extension of structure group the rational principal H -bundle $P(H)$ is also semistable.*

Proof: By the choice of P and by the restriction theorem of Mehta and Ramanathan, we may choose a high degree general smooth complete intersection curve C of type (m) , with $C \subset U$ such that $E|_C$ is polystable of degree zero. Let a denote the product $a_1 \dots a_{d-1}$ in the notation of (3.1).

Thus by Prop 3.11, the associated H -bundle $P(H)|_C$ is a semistable principal bundle on C .

We *claim* that this implies that $P(H)$ is itself a semistable principal H -bundle. For, if $Q \subset H$ is a parabolic subgroup and χ a dominant character of Q , $P(H)_Q$ a Q -bundle obtained from a reduction of structure group to Q , note that

$$\deg P(H)_Q(\chi) \cdot a^m = \deg P(H)_Q(\chi)|_C$$

where $P(H)_Q(\chi)$ denotes the line bundle associated to the character χ . Since $m \gg 0$ it follows by the semistability of $P(H)|_C$ that $\deg P(H)_Q(\chi)|_C > 0$ and hence $\deg P(H)_Q(\chi) > 0$, i.e $P(H)$ is semistable.

q.e.d

Proof.(of Theorem 4.1) Let the semistable principal H_K -quasibundle P_K arise out of the faithful representation $H \subset G = GL(V)$. In other words, there is a pair $(\mathcal{E}_K, \sigma_K)$ of semistable torsion-free sheaf $\mathcal{E}_K$ on X_K and section σ_K . By Prop 2.13, we have a semistable extension $\mathcal{E}_A$ of $\mathcal{E}_K$ (possibly by going to a finite extension L/K such that $\mathcal{E}_p$ is polystable of degree 0. Let $U = U(\mathcal{E}_p)$. Then as we have seen earlier, the entire family $\mathcal{E}_A$, when restricted to U_A is locally free. Now we are in the setting of Prop 3.7.

Moreover, the fibre of E_B over the closed point is indeed **semistable**. To see this, observe firstly that it comes as the extension of structure group of E'_p by the map $\psi_k : H'_k \longrightarrow H_k$. Recall (Proposition 3.8) that E'_p is the *semistable* H'_k -bundle obtained as the reduction of structure group of the polystable vector bundle $E(V_A)_p$ and the semistability of the extended H_k -bundle follows by Lemma 4.4.

By Proposition 3.7 we have a *rational* H'_A -bundle P'_A on U_A , which extends the H_K -bundle upto isomorphism. Further, by Proposition 4.3, by going to the extension L/K we have a morphism of B -group schemes $\psi_B : H'_B \longrightarrow H_B$ which is an isomorphism over L . Therefore, one can extend the structure group of the bundle P'_B to obtain a rational H_B -bundle P_B which extends the H_K -bundle P_K . By using the faithful representation $H \subset GL(V)$, we get a family of vector bundles $P_B(V)$ on U_B .

Note that the rational H'_K -bundle P_K is the restriction to U_K of the principal H_K -quasibundle $(\mathcal{E}_K, \sigma_K)$.

In other words, we have an isomorphism $P_L(V) \simeq \mathcal{E}_L$ on U_L .

Now we apply a result from Langton's paper ([22, Prop 6]) to conclude that the bundle $P_B(V)$ on U_B in fact extends to a torsion-free sheaf $\mathcal{E}_B$ on X_B . Further, the reduction section σ_L (defined on X_L has extended to a section σ_B on U_B (given by the principal H_B -bundle P_B). We can view the reduction datum as follows:

$$\sigma_B : (U_B \cup X_L) \longrightarrow (S_{\mathcal{E}_B})//H_B$$

(which by the extension property, agrees on the intersection). The complement of $U_B \cup X_L$ in X_B is a subset of points of codimension ≥ 3 , and one can embed the affine scheme $(S_{\mathcal{E}_B})//H_B$ as a *closed subscheme* of a vector bundle over X_B .

This implies that the section σ_B extends to a section $\sigma_B : X_B \longrightarrow (S_{\mathcal{E}_B})//H_B$. In other words, the pair $(\mathcal{E}_K, \sigma_K)$ has been semistably extended (upto isomorphism) to a pair $(\mathcal{E}_B, \sigma_B)$, i.e, the principal H_K -quasibundle P_K has been semistably extended (upto isomorphism) to principal H_B -quasibundle P_B , possibly after going to a finite extension B/A . This completes the proof of the Theorem 4.1.

q.e.d

For purposes of applications later we isolate the following easier half of the Mehta–Ramanathan restriction theorem:

LEMMA 4.5. *Let P be an H -quasibundle. Then P is μ -polystable (resp stable) if $P|_C$ is so for a general high degree smooth complete intersection curve $C \subset X$.*

Proof: The proof is along the lines of the first part of the proof of Prop 4.4. For more general results along similar lines cf. [7], whose methods work easily enough for rational principal bundles and hence for quasibundles.

5. CONSTRUCTION OF THE MODULI SPACE OF BUNDLES OVER SURFACES

From now onwards X will be a *smooth projective surface*. The aim of this section is to give an algebro-geometric construction of the Donaldson–Uhlenbeck compactification for the moduli space of semistable principal H -bundles. This, in particular also describes geometrically the Yang-Mills

compactification of the moduli space of ASD connections on principal bundles with arbitrary structure groups. The method of proof is along the lines of the proof of J.Li (cf. [24]) and the methods in the paper of Le Potier ([23]) (cf. also [19, Chapter 8]).

5.0.1. *Double duals for quasibundles.* If $P|_U$ is a rational H -bundle, where U is a *big* open subset of X , then by a theorem of Colliot-Thélène and Sansuc([10, Theorem 6.7]) there is a principal bundle (unique upto isomorphism) which extends $P|_U$. This extension is therefore also associated to each quasibundle P as well.

DEFINITION 5.1. (*Double duals*) We call this extended principal bundle the double dual of the quasibundle P and denote it by P^{**} . This plays the role of the *double dual* of torsion free sheaves.

Remark 5.2. If P is a μ -semistable (resp. polystable, stable) H -quasibundle, then since these definitions involve only *big* open subsets, the corresponding double dual P^{**} is a μ -semistable (resp. polystable, stable) principal H -bundle.

5.0.2. *On determinant line bundles.* Recall that if $\mathcal{F}$ is a flat family of coherent sheaves on X parametrised by a scheme S , then $\mathcal{F}$ defines an element $[\mathcal{F}] \in K^0(S \times X)$, the Grothendieck group of $S \times X$ generated by locally free sheaves. We may then define the homomorphism from the Grothendieck group of coherent sheaves on X given by:

$$\lambda_{\mathcal{F}} : K(X) \longrightarrow \text{Pic}(S).$$

This has a collection of functorial properties for which we refer to ([19] page 179). For every class $u \in K(X)$ we denote the associated class of the line bundle by $\lambda_{\mathcal{F}}(u)$. Fix a class $c \in K(X)$ with rank r and Chern classes $c_1 = \mathcal{O}_X$ and c_2 , the very ample divisor Θ on X and a base point $x \in X$. Then there is a natural choice of a class $u_1(c) \in K(X)$ defined in terms of these fixed data (cf. page 184 [19]).

5.0.3. *The parameter space for quasibundles.* We first briefly recall the construction of the parameter space of μ -semistable torsion-free sheaves over the surface X . Let $c \in K(X)$ be as above. Then one knows that the set $\mathcal{S}$ of isomorphism classes of μ -semistable torsion-free sheaves of class c with trivial determinant and fixed Hilbert polynomial P is bounded and hence for a $m \gg 0$ we can realise them as points of a suitable quot scheme. Fix such an m . Let W be a complex vector space of dimension $P(m)$.

Let $R^{\mu ss} \subset \text{Quot}(W, P)$ be the locally closed subscheme of all quotients $[q : W \otimes \mathcal{O}_X(-m) \longrightarrow \mathcal{E}]$ such that $\mathcal{E}$ is μ -semistable of rank r with trivial determinant and second Chern class c_2 , with Hilbert polynomial P and such that q induces an isomorphism $M \simeq H^0(\mathcal{E}(m))$.

There is a natural action of $\mathcal{G} = SL(M)$ on $R^{\mu ss}$ and we have the universal quotient $q_Q : \pi_X^* \mathcal{O}_X(-m) \otimes W \longrightarrow \mathcal{E}_Q$.

Returning to our setting, let $\rho : H \hookrightarrow G$ be a faithful representation of H in $G = SL(V)$ where $\dim(V) = r$. We shall use the following notation when we wish to stress the fact that the quot scheme is parametrising semistable principal quasibundles with structure group G rather than torsion-free sheaves of rank r :

$$R_G := R^{\mu ss}$$

Denote the universal quotient on $X \times R_G$ by $q_{R_G} : \pi_X^* \mathcal{O}_X(-m) \otimes W \longrightarrow \mathcal{E}_G$.

Recall our notation of §2 and the notion of quasibundles with respect to the representation ρ . These are obtained from the affine X -scheme $S_{\mathcal{E}}$ together with a *generalised reduction* datum $\sigma : X \longrightarrow S_{\mathcal{E}}//H$. The pair $(\mathcal{E}, \sigma)$ is a principal quasibundle with structure group H . We remark that this reduction datum σ can be equivalently viewed as giving an $\mathcal{O}_X$ -algebra morphism $\tau : (\text{Sym}^*(\mathcal{E} \otimes V))^H \longrightarrow \mathcal{O}_X$. The map τ which comes from a *genuine* generalised reduction σ is not simply the projection onto $\mathcal{O}_X$. It is obtained by dualising the map σ , where $\sigma|_U$ is a reduction of structure group of the principal bundle (associated to $\mathcal{E}|_U$) to H on a *large* open subset $U = U(\mathcal{E})$.

Let $R_H(\rho)$ be the scheme which parametrises pairs (q, τ) where $q \in R_G$ and τ is a reduction datum giving a *quasibundle with structure group H* and let $R_H(\rho)^o$ be the open subset of pairs (q, τ) such that τ defines a *principal bundle* with structure group H .

The existence of total families for principal quasibundles follows by the general theory of Hilbert schemes and is shown in ([35]); in Schmitt's language these will be parametrising what he terms *honest singular bundles*. We have been somewhat loose in defining the parameter space but we refer the reader to Schmitt's paper (cf. [35, Section 6.7] for details).

The scheme $R_H(\rho)$ is an R_G -scheme and the natural map $f_{\rho} : R_H(\rho) \longrightarrow R_G$ is the one induced by ρ . Therefore, since we have already fixed the Chern classes of the torsion free sheaves in R_G , by the general theory of characteristic classes for principal bundles, the characteristic classes of the principal bundles in the open subscheme $R_H(\rho)^o \subset R_H(\rho)$ will also be canonically fixed. This follows by the basic result of Borel–Hirzebruch which relates the characteristic classes of principal bundles with the Chern classes of associated vector bundles. For a nice exposition and explicit results cf. [6, Prop 3.2].

For simplicity we will denote this entire datum by $\mathbf{c}$ (we will return to this in 6.2.1).

Remark 5.3. It is immediate that the $\mathcal{G}$ -action on R_G lifts to an action on $R_H(\rho)$.

By the universal property of the scheme $R_H(\rho)$ it follows that there exists a tautological family on $X \times R_H(\rho)$. Let $\mathcal{P}$ denote this $R_H(\rho)$ -flat family of H -quasibundles on X associated to the faithful representation ρ .

Then by its definition, there exists an $R_H(\rho)$ -flat family of torsion free sheaves $\mathcal{P}(\rho) = \mathcal{F}$ on $X \times R_H(\rho)$. This is given precisely by the family $(id_X \times f_\rho)^*(\mathcal{E}_G)$ obtained from the family of semistable torsion free sheaves on $X \times R_G$. As remarked above, there is $\mathcal{G}$ -action on $R_H(\rho)$ such that the family $\mathcal{F}$ carries a linearisation with respect to the $\mathcal{G}$ -action.

DEFINITION 5.4. Let $\mathcal{P}(\rho) = \mathcal{F}$ be the induced family of torsion free sheaves on X parametrised by $R_H(\rho)$. Fixing the class $c \in K(X)$ as seen above, we have a canonical choice of $u_1(c) \in K(X)$. We then have the *determinant line bundles* $\mathcal{L} := \lambda_{\mathcal{F}}(u_1(c))$ on $R_H(\rho)$ induced by the family $\mathcal{F}$.

Notation Since we have emphasized the role of the representation ρ , we will henceforth denote the total space $R_H(\rho)$ simply by R_H .

Then we have the following:

LEMMA 5.5. ([19, Lemma 8.2.4])

1. If $s \in S$ is a point such that $\mathcal{P}_s|_C$ is μ -semistable. Then there exists an integer $n > 0$, and $\mathcal{G}$ -invariant sections $\bar{\sigma}$ in $\lambda_F(u_1(c))^n$ such that $\bar{\sigma}(s) \neq 0$.
2. Let s_1 and s_2 be two points in S such that either $\mathcal{P}_{s_1}|_C$ and $\mathcal{P}_{s_2}|_C$ are both semistable but not S -equivalent (in the sense of Seshadri) or one of them is semistable and the other is not. Then, in either case, there are $\mathcal{G}$ -invariant sections in some tensor power of $\lambda_F(u_1(c))$ that separate s_1 and s_2 .

Proof: Since the proof follows the general line as given in the text ([19]), we content ourselves by giving the main steps in the argument and refer the reader to [19] for more details.

We need to relate the “determinant line bundle” on R_H to its pull-back from the corresponding total family Q_C of principal G -bundles on a high degree curve $C \subset X$.

Let C be a general smooth curve in X of high degree $a \gg 0$. Let $i : C \hookrightarrow X$. For the class $c \in K(X)$, its restriction, to $c|_C := i^*(c)$ in $K(C)$, is completely determined by its rank r since we have assumed that all our sheaves have trivial determinant. Let P' be the Hilbert polynomial determined on C by this restriction. Then for a large positive integer m' we consider the induced “quot scheme” of quotients $[q : W' \otimes \mathcal{O}_C(-m') \rightarrow E]$. Let us denote this quot scheme by Q_C . Here W' is a complex vector space of dimension $P'(m')$. Let Q_C^{ss} be the quotients which give semistable bundles on C and let $q_C : W' \otimes \pi_C^*(\mathcal{O}_C(-m')) \rightarrow \mathcal{E}_C$ be the universal quotient on $X \times Q_C$ with *determinant line bundle* $\mathcal{L}'_0$ on Q_C obtained from this family. We then have the following properties on *separating points* in Q_C by sections of the determinant line bundle $\mathcal{L}'_0$:

- (a) All $SL(P'(m'))$ -semistable points (in the GIT sense) in Q_C are precisely the bundles E which are semistable. Further, there exists an

integer ν and an $SL(P'(m'))$ -invariant section σ of $\mathcal{L}'_0{}^\nu$ such that $\sigma([q]) \neq 0$, where $q \in Q_C^{ss}$ corresponds to the bundle E .

- (b) Two points q_i , $i = 1, 2$ in Q_C are separated by invariant sections in some tensor power of $\mathcal{L}'_0$ if and only if either both are semistable but not S -equivalent or one of them is semistable and the other is not.

We now work with R_H and the family $\mathcal{F}$ on $X \times R_H$. Let us denote the restriction of $\mathcal{F}$ to $C \times R_H$ by $\mathcal{F}_C$.

Now by increasing m' if necessary, we can make sure that:

- The restrictions $\mathcal{F}_s|_C$ are points of Q_C .
- The sheaf $p_*\mathcal{F}_C(m')$ is a locally free $\mathcal{O}_{R_H}$ -module of rank $P'(m')$, where $p : C \times S \rightarrow S$ is the projection.

Let S_H be the associated projective frame bundle and let $\eta : S_H \rightarrow R_H$ be the natural map. It parametrises a quotient

$$\mathcal{O}_{S_H} \otimes \pi_C^* \mathcal{O}_C(-m') \otimes W' \rightarrow \eta^* \mathcal{F}_C \otimes \mathcal{O}_\eta(1).$$

This induces a $SL(P'(m'))$ -invariant morphism

$$\Phi_{\mathcal{F}_C} : S_H \rightarrow Q_C$$

We also have the $\mathcal{G}$ -action on $\mathcal{F}$ and via this action we get a $\mathcal{G}$ -action on S_H which commutes with the $SL(P'(m'))$ -action such that both $\eta : S_H \rightarrow R_H$ and $\Phi_{\mathcal{F}_C}$ are equivariant for the $\mathcal{G} \times SL(P'(m'))$ -actions.

One then shows by a degree computation that:

$$\Phi_{\mathcal{F}_C}^*(\mathcal{L}'_0)^{\deg C} \simeq \eta^* \mathcal{L}^{a^2 \deg(X)} \quad (\#)$$

(The canonical *det* line bundle on R_H depends only on the choice of a polarisation on X and the relation $\#$ is independent of $C \in |a\Theta|$. cf. [19] pp 187-188)

Since $S_H \rightarrow R_H$ is a principal $PGL(P'(m'))$ -bundle, a section invariant under $SL(P'(m'))$ descends to give a $\mathcal{G}$ -invariant section. Thus, for any ν we thus get a linear map:

$$s_{\mathcal{F}} : H^0(Q_C, (\mathcal{L}'_0)^{\nu \deg(C)})^{SL(P'(m'))} \rightarrow H^0(R_H, \mathcal{L}^{\nu a^2 \deg(X)})^{\mathcal{G}}$$

To conclude the proof of the Lemma we first observe that the semistability of the quasibundle $\mathcal{P}_s$ is equivalent to the μ -semistability of the torsion-free sheaf $\mathcal{F}_s$ for each $s \in R_H$. Hence by using section of powers of the line bundle $\mathcal{L}'_0$ on Q_C and via the map $s_{\mathcal{F}}$ we can indeed separate the restrictions of quasibundles in R_H to C as well.

q.e.d

Remark 5.6. By the Mehta–Ramanathan theorem, if the degree of the curve C is large enough, then the restriction of a μ -semistable torsion-free sheaf on X to the curve C is semistable. But the same curve C may not work for all the bundles in R_H . Hence we need to work with the whole of Q_C .

We have the following immediate corollary from the first part of Lemma 5.5:

COROLLARY 5.7. *There exists an integer $\nu > 0$ such that the line bundle $\mathcal{L}^\nu$ on R_H is generated by $\mathcal{G}$ -invariant global sections i.e $\mathcal{L}$ is $\mathcal{G}$ -semi-ample.*

Since R_H is a quasi-projective scheme and since $\mathcal{L}$ is $\mathcal{G}$ -semi-ample, there exists a finite dimensional vector space $J \subset J_\nu := H^0(R_H, \mathcal{L}^\nu)^\mathcal{G}$ that generates $\mathcal{L}^\nu$. Note that there is nothing canonical in the choice of J .

Let morphism $\phi_J : R_H \rightarrow \mathbb{P}(J)$ be the induced $\mathcal{G}$ -invariant morphism defined by the sections in J . Because of nonuniqueness of J , each choice of subspace of invariant sections gives rise to a different map $\phi_{J'}$ to a different projective space $\mathbb{P}(J')$.

DEFINITION 5.8. We denote by M_J the **schematic image** $\phi_J(R_H)$ with the **canonical reduced scheme structure**.

Remark 5.9. By the following result which may be titled $\mathcal{G}$ -properness, the variety M_J is *proper* and hence because of its quasi-projectivity it is a projective variety. We note that we use the term *variety* in a more general sense of an *reduced algebraic scheme of finite type* which need not be irreducible. So in what follows we will be working with the $\mathbb{C}$ -valued points of M_J .

PROPOSITION 5.10. *If T is a separated scheme of finite type over k , and if $\phi : R_H \rightarrow T$ is an $\mathcal{G}$ invariant morphism then image of ϕ is proper over k .*

Proof: This is an immediate consequence of Theorem 4.1 and [19, Prop 8.2.6].

q.e.d

Let J_ν denote the vector space $H^0(R_H, \mathcal{L}^\nu)^\mathcal{G}$, $\nu \in \mathbb{Z}^+$; and Let $J \subset J_\nu$ be a finite dimensional vector space which generates $\mathcal{L}^\nu$.

For any $d \geq 1$, let J^d be the image of the canonical multiplication map $f_d : J \otimes \cdots \otimes J (d - \text{fold}) \rightarrow J_{d\nu}$; in particular, $J^1 = J$.

Let J' be any finite dimensional vector subspace of $J_{d\nu}$ containing J^d . Then clearly the line bundle $\mathcal{L}^{d\nu}$ is also globally generated by $\mathcal{G}$ -invariant sections coming from the subspace J' and this is so for any $d \geq 0$.

So we have inclusions $J \hookrightarrow J^d \hookrightarrow J'$, and hence a commutative diagram

$$\begin{array}{ccc} M_{J'} & \xrightarrow{\pi_{J'/J}} & M_J \\ \uparrow \phi_{J'} & \nearrow \phi_J & \\ R_H & & \end{array}$$

Since M_J and $M_{J'}$ are both projective, the map $\pi_{J'/J}$ is a finite map. So if we fix a J as above, we get an inverse system (indexed by the $d \geq 1$) of projective varieties $(M_{J'}, \pi_{J'/J})$ and dominated by the finite type scheme R_H .

$$\begin{array}{ccccc}
& & R_H & & \\
& \swarrow \phi_{J'} & \downarrow & \searrow \phi_J & \\
\cdots & \longrightarrow & M_{J'} & \xrightarrow{\pi_{J'/J}} & M_J
\end{array}$$

Hence the inverse limit of the system $(M_{J'}, \phi_*)$ is in fact one of the $M_{J'}$'s where J' is a finite dimensional subspace of $H^0(R_H, \mathcal{L}^n)^{\mathcal{G}}$ which generates $\mathcal{L}^n$.

DEFINITION 5.11. We denote this *inverse limit variety* by $M_H(\rho)$ and let $\pi : R_H \rightarrow M_H(\rho)$ be the canonical morphism induced by the invariant sections coming from the subspace J_0 associated to the inverse limit.

We summarise the above discussion in following theorem:

THEOREM 5.12. *Then the reduced projective scheme $M_H(\rho)$ parametrises equivalence classes of μ -semistable quasibundles with structure group H . There is a natural morphism $q : R_H \rightarrow M_H(\rho)$. Furthermore, the representation $\rho : H \rightarrow G = SL(V)$ induces a natural morphism $\bar{f}_\rho : M_H(\rho) \rightarrow M_G$, where $M_G := M_{SL(V)}$ is the moduli space of the μ -semistable torsion free sheaves with trivial determinant and fixed c_2 .*

Proof: The existence of the morphism $\bar{f}_\rho : M_H(\rho) \rightarrow M_G$ follows by the *naturality* of the moduli space M_G by virtue of the existence of the family of semistable torsion free sheaves $\mathcal{F}$ on $R_H(\rho)$.

q.e.d

REMARK 5.13. The strategy is somewhat similar to that of [3] but unlike [3], there are no GIT quotients involved here.

REMARK 5.14. Note that this is not a categorical quotient since $\mathcal{L}$ is not ample and is only *semi-ample* (Cor 5.7), i.e some power of $\mathcal{L}$ is generated by sections.

5.1. **Towards the description of the moduli space.** In order to get a better understanding of the geometry of the moduli space $M_H(\rho)$ we need the following lemmas:

LEMMA 5.15. *Let P_1 and P_2 be two polystable principal H -bundles on X . Let $a \gg 0$ and $C \in |a\Theta|$ be a general smooth curve. Then $P_1 \simeq P_2$ if and only if $P_1|_C \simeq P_2|_C$.*

Proof: Consider the principal $H \times H$ -bundle $P_1 \times_X P_2$. Let us denote this bundle by E .

By ([37], p 19), to give an isomorphism between P_1 and P_2 is equivalent to giving a *reduction of structure group* of E to the diagonal embedding $\Delta \subset (H \times H)$.

By the Chevalley semi-invariant theorem, we can embed $(H \times H)/\Delta \subset W$ in a $H \times H$ -module W as a closed orbit.

Thus a section of $E(\frac{H \times H}{\Delta})$ is a section of the vector bundle $E(W)$ which lies in $E(\frac{H \times H}{\Delta})$.

Claim: The principal bundle E is also polystable of degree 0. Let us assume this claim and complete the proof.

By the usual Enriques–Severi lemma for a general high degree curve C , the restriction map:

$$H^0(X, E(W)) \longrightarrow H^0(C, E(W)|_C) \quad (\&)$$

is *surjective*. Further, by assumption, for high degree curves we know that $P_1|_C \simeq P_2|_C$. Hence, by our discussion above we have a reduction of structure group of $E|_C$ to Δ . i.e a section:

$$\sigma_C : C \longrightarrow E(\frac{H \times H}{\Delta})|_C \hookrightarrow E(W)|_C$$

By (&) above, there exists a lift of σ_C to a section $\sigma : X \longrightarrow E(W)$. We need to show that the image of σ lies in $E(\frac{H \times H}{\Delta})$.

One knows that E is polystable of degree 0, by the *Claim* above. Thus we are in the setting of Prop 3.13. Thus we can get a big open subset $U \subset X$ such that the image of the section $\sigma(U) \subset E(\frac{H \times H}{\Delta})$.

Now since $\frac{H \times H}{\Delta} \subset W$ is a closed embedding, it follows that the entire image $\sigma(X) \subset E(\frac{H \times H}{\Delta})$. This gives the required reduction of structure group to Δ . To complete the proof we need to prove the claim.

To see this we again use the easier half of the Mehta–Ramanathan restriction theorem (i.e Lemma 4.5). Thus to show that E is polystable of degree 0 we need to show that for a general high degree curve C , the restriction $E|_C$ is polystable of degree zero.

Since $P_1|_C$ and $P_2|_C$ are polystable of degree 0 (by the restriction theorem again), it follows by Ramanathan’s theorem ([31]) (the Narasimhan–Seshadri theorem for principal bundles on curves) that there exists representations $\rho_i : \pi_1(C) \longrightarrow K$ $i = 1, 2$, K a maximal compact subgroup of H , such that $P_i|_C$ is the H -bundle associated to ρ_i , for $i = 1, 2$.

It is easy to see that the bundle $E|_C$ is the bundle associated to the representation $\rho_1 \times \rho_2$ ([31] p 146) and the polystability follows by Ramanathan’s theorem again. This completes the proof of the Lemma.

q.e.d

5.1.1. Associated graded of a semistable quasibundle. Let $P \longrightarrow X$ be an H -quasibundle. Let $\rho : H \hookrightarrow GL(V)$ be the accompanying faithful representation and $\mathcal{E} = P(\rho)$ the associated torsion-free sheaf. The sheaf $\mathcal{E}$ possesses a Jordan–Holder filtration $J^\bullet$ by saturated subsheaves and we may take the associated graded sheaf $gr_{J^\bullet}(\mathcal{E})$. For two different filtrations $J_i^\bullet$ we have

an isomorphism of vector bundles $(gr_{j_1}(\mathcal{E}))^{**} \simeq (gr_{j_2}(\mathcal{E}))^{**}$. By an abuse of notation we write $gr^\mu(\mathcal{E})$ to denote an associated graded torsion-free sheaf.

Let $\mathbb{A} = \mathbb{A}_{\mathbb{C}}^1$. Recall that there exists a family $\{\mathcal{E}_t\}_{t \in \mathbb{A}}$ of torsion-free sheaves such that $\mathcal{E}_t \simeq \mathcal{E}$ for $t \neq 0$ and $\mathcal{E}_0 \simeq gr^\mu(\mathcal{E})$.

By the *Semistable reduction Theorem 4.1 (and its proof)* we have an H_A -quasibundle (where $Spec A = \mathbb{A}_1 \subset \mathbb{A}$ an open subset which contains 0). Denote this H_A -quasibundle by $\{P_t\}_{t \in \mathbb{A}_1}$. Further, this family has the property that $P_t \simeq P$ for $t \neq 0$ and there is a big open set U and a rational H' -bundle P'_0 on U , such that $P_0|_U$ is obtained from P'_0 by an extension of structure group via $\psi : H' \rightarrow H$ (recall that H' could be a non-reductive group; cf. Prop 4.3 and Lemma 4.4).

Also from the proof (Theorem 3.7) one knows that $P'_0(\rho) \simeq \mathcal{E}_0|_U$ which is *polystable*.

Claim: The quasibundle P_0 is also polystable.

By Lemma 4.5 it is enough to show that $P_0|_C$ is polystable, where C is a high degree curve contained in the big open subset U , where P_0 is a principal bundle. Further, $P'_0|_C$ is a bundle which is *flat* i.e it comes from a representation $\chi_1 : \pi_1(C) \rightarrow H'$ (see Prop 3.8). Moreover, $P'_0(\rho)|_C \simeq \mathcal{E}_0|_C$ which is *polystable* by the Mehta–Ramanathan theorem, since $\mathcal{E}_0|_U$ is polystable. In other words, under our definition of polystability of principal bundles with non-reductive structure groups (i.e Def 3.9), the bundle $P'_0|_C$ is *polystable*.

Again $P_0|_C$ comes from $P'_0|_C$ by extension of structure group $f : H' \rightarrow H$. Therefore, since $P'_0|_C$ is polystable by Prop 3.11, it follows that $P_0|_C$ is *polystable*. This proves the claim.

DEFINITION 5.16. Let P be a semistable H -quasibundle. If there exists a family $\{P_t\}_{t \in Spec A}$, with A a complete discrete valuation ring, such that $P_t \simeq P$, for $t \neq 0$ and P_0 polystable. Then we call P_0 **an** associated graded quasibundle of P .

This is uniquely defined upto *double duals* in the following sense:

LEMMA 5.17. *Let P be an H -quasibundle. Then if P_{s_0} and P_{t_0} are two choices of polystable limits (as above), then $P_{s_0}^{**} \simeq P_{t_0}^{**}$.*

Proof: Let P_S and P_T be two families of quasibundles such that at the closed points $s_0 \in S$ and $t_0 \in T$ the quasibundles P_{s_0} and P_{t_0} are polystable and the generic fibres in either family is isomorphic to P .

Consider the open subset $U \subset X$ where both P_{s_0} and P_{t_0} are locally free. Then since S and T are spectra of discrete valuation rings, the families P_S and P_T are locally free on $U \times S$ and $U \times T$ respectively. Using the Mehta–Ramanathan theorem choose a general high degree curve $C \subset U$ so that the restrictions $P_{s_0}|_C$ and $P_{t_0}|_C$ are polystable. Since $C \subset U$ it follows that both $P_{s_0}|_C$ and $P_{t_0}|_C$ are polystable limits of $P|_C$ (which is also semistable by openness of semistability). But the associated graded of a semistable

principal bundle on a smooth projective curve is uniquely defined (cf. [32]). Hence,

$$P_{s_0}|_C \simeq P_{t_0}|_C.$$

Now we apply Lemma 5.15 to the double dual principal bundles on X and we are done.

Remark 5.18. If $P(\rho) = \mathcal{E}$, then for each choice of associated H -quasibundle $gr^\mu(P) = P_0$ defined above, we see that $P_0^{**}(\rho) \simeq \mathcal{E}_0^{**}$.

6. THE GEOMETRY OF THE MODULI OF H -BUNDLES

From this section onwards H is a simple algebraic group. In this section we study the points of the moduli space $\overline{M}_H^0$ intrinsically as well as in relationship with its image points in the moduli space M_G .

6.1. Cycles associated to quasibundles in $\overline{M}_H^0$. Let P be a semistable H -quasibundle in $\overline{M}_H^0$. Therefore there exists a family $\{P_t\}_{t \in \text{Spec } A}$, with A a complete discrete valuation ring, such that P_t is a semistable principal H -bundle, for $t \neq 0$ and $P_0 \simeq P$.

Our aim is to associate a cycle $Z_P \in S^l(X)$ of degree l to the quasibundle P in an intrinsic manner. By this we mean that the pair (P^{**}, Z_P) in the compactified moduli space is independent of the quasibundle (and hence the representation ρ as well). Towards this we work with the chosen representation ρ used in defining the quasibundle P .

6.1.1. The Dynkin index of the representation ρ . We recall the notion of Dynkin index ([13], [21], [2]) and some basic results from these sources which will play a key role in what follows.

DEFINITION 6.1. Let θ be a the highest root in $\mathfrak{h}$ the Lie algebra of H and $\mathfrak{sl}(\theta)$ the 3-dimensional sub-algebra of $\mathfrak{h}$ associated to it. Decompose the H -module V into $\mathfrak{sl}(\theta)$ -modules as $\oplus V_i = V$. Let $\dim(V_i) = m_i$. Then we can define the Dynkin index m_ρ of the H -module V as follows:

$$m_\rho = \sum_i \binom{m_i + 1}{3}$$

Remark 6.2. The Dynkin index of a simple subgroup H of a simple Lie group G is usually defined as the ratio of the invariant inner product on $\text{Lie}(H)$ to the invariant inner product of $\text{Lie}(G)$ where the inner products are normalised to make the length of the highest root 2 ([2, page 455]).

Example 6.3.

- (a) Let $\rho_m : SL(2) \longrightarrow SL(V)$ be the standard irreducible representation $V = S^m(W)$ where $\dim(W) = 2$. The $m_{\rho_m} = \binom{m+2}{3}$.
- (b) Let $\rho : SL(2) \longrightarrow SL(V)$ be the inclusion obtained by identifying $SL(2)$ with the 3-dimensional group given by the highest root θ in $\mathfrak{sl}(V)$. Then $m_\rho = 1$.

- (c) If ρ is the adjoint representation of H then the index is $2h$, where h is the dual Coxeter number of H .

The following is the list of index 1 subgroups given by Dynkin ([13]). We reproduce the list from [2]:

TABLE 1
$S(U(n) \times U(1)) \subset SU(n+1)$
$Spin(n) \subset Spin(n+1)$
$Sp(n) \times Sp(1) \subset Sp(n+1)$
$SU(3) \subset G_2$
$Spin(9) \subset F_4$
$F_4 \subset E_6$
$E_6 \times U(1) \subset E_7$
$E_7 \times SU(2) \subset E_8$

We now quote from [21] (see also [2, page 455]);

PROPOSITION 6.4. *Let $\rho_* : \pi_3(H) \longrightarrow \pi_3(SL(V))$. Then it is given by “multiplication” by m_ρ .*

Remark 6.5. This in particular proves that any index 1 inclusion $SU(2) \subset SU(n)$ preserves the second Chern class of a principal $SU(2)$ -bundle when we extend structure group to $SU(n)$ since $\pi_3(G) \simeq \mathbb{Z}$ classifies principal G -bundles on a real 4-sphere S^4 and the second Chern class of the bundle classifies it. This also implies that any $SU(n)$ -bundle E on S^4 is obtained as the extension of structure group from $SU(2)$. In particular, the the rank n vector bundle underlying E (topologically) splits as a direct sum of a rank 2 bundle and direct sum of $n-2$ trivial line bundles. Thus if the vector bundle underlying E has a nontrivial section s then the section comes from a section of the rank 2 subbundle. Hence the “zero-scheme” $Z(s)$ is represented by $c_2(E)$.

For a point $x \in X$, let $B_\epsilon(x)$ be an analytic ball of radius ϵ around x . Let E be a vector bundle on X of rank r and let $E|_{B_\epsilon(x)}$ be its restriction to $B_\epsilon(x)$. Suppose further that it is trivial on the boundary $\partial B_\epsilon(x)$. Let us denote by $E|_{B_\epsilon(x)}/E|_{\partial B_\epsilon(x)}$ the complex vector bundle on the real four sphere $S^4 \simeq B_\epsilon(x)/\partial B_\epsilon(x)$ obtained by identifying $E|_{\partial B_\epsilon(x)}$ to $\mathbb{C}^{\oplus r}$ using the trivialisation.

We then have the following result from [24] and [28] extended for rank r bundles. Li and Morgan show it for rank 2 case but the proofs generalise to the higher rank case.

PROPOSITION 6.6. *Let $\mathcal{E}_\mathbb{A}$ be a family of (analytic) torsion-free sheaves of rank r over $B \times \mathbb{A}$ flat over $\mathbb{A}$, where B is a closed ball in the surface X and $\mathbb{A}$ is the affine line. Further let $\mathcal{E}_\mathbb{A}$ be locally free on $B \times \mathbb{A} - 0$ and $\mathcal{E}_0$ be torsion-free with singularity at the origin $0 \in B$. Let $B_{\epsilon,u} \subset B \times \{u\}$ be the 2-dimensional ϵ -ball centered at $(0, u)$ and let $\mathcal{E}_u$ be the restriction of $\mathcal{E}$ to $B_{\epsilon,u}$. Then the smooth trivialisation of $\mathcal{E}_0|_{\partial B_{\epsilon,u}}$ which is induced from a trivialisation of $(\mathcal{E}_0|_{B_{\epsilon,u}})^{**}$ induces a family of trivialisations $\beta_u :$*

$\mathbb{C}^{\oplus r} \times \partial B_{\epsilon,u} \simeq \mathcal{E}_{u|_{\partial B_{\epsilon,u}}}$ which depends smoothly on u with $u \in \mathbb{A}$ being ϵ -small. Furthermore, we also have on $S^4 \simeq B_{\epsilon,u}/\partial B_{\epsilon,u}$ the following:

$$c_2(\mathcal{E}_{u|_{B_{\epsilon,u}}}/\mathcal{E}_{u|_{\partial B_{\epsilon,u}}}) = \text{length}(\mathcal{E}_0^{**}/\mathcal{E}_0)_0$$

Proof: The only new ingredients needed to generalise Li's arguments are:

- Extension of [24, Prop 6.4] to higher ranks which has been done by Gieseker and Li [17].
- When the length $l = 1$ then the length l can be realised as the second Chern class of the rank r bundle on S^4 . This is done in Li in the rank 2 case by getting a section whose zero-scheme has length 1 and is represented by the c_2 . Since we are on S^4 this argument goes through for higher rank case as well as we have seen in Remark 6.5.

q.e.d

Let $P_{t_0} \in \overline{M_H^0}$ be a polystable H -quasibundle and P_T be a family of semistable principal H -bundles with P_{t_0} as limit. Let $\mathcal{E}_T = P_T(\rho)$. Let $Z(\mathcal{E}_{t_0})$ be the cycle associated to the torsion-free sheaf $\mathcal{E}_{t_0}$. Recall that $Z(\mathcal{E}_{t_0}) \in S^l(X)$ where $l = c_2(\mathcal{E}_{t_0}^{**}) - c_2(\mathcal{E}_{t_0})$. Further the cycle is given by:

$$Z(\mathcal{E}_{t_0}) := \sum_{x \in X} \text{length}(\mathcal{E}_{t_0}^{**}/\mathcal{E}_{t_0})_x \cdot x$$

THEOREM 6.7. *Let the notations be as above. Let $x \in \text{Supp}(Z(\mathcal{E}_{t_0}))$. Then the number $\text{length}(\mathcal{E}_{t_0}^{**}/\mathcal{E}_{t_0})_x$ is a multiple of the Dynkin index m_ρ . In particular, the total degree l of the cycle $Z(\mathcal{E}_{t_0})$ is also a multiple of m_ρ .*

Proof: Fix a point $x \in \text{Supp}(Z(\mathcal{E}_{t_0}))$ and choose an ball B around it and identify x with $0 \in B$. Let $u \in \mathbb{A}$ is the point in a small disk in $\mathbb{A}$ corresponding to $t_0 \in T$.

By restricting $\mathcal{E}_T$ to $B \times \mathbb{A}$, in the notation of Prop 6.6, we have a family $\mathcal{E}_{\mathbb{A}}$ satisfying the properties given there. Furthermore, we also have the extra datum that the bundles $\mathcal{E}_t$ for $t \neq t_0$ have reduction of structure group to principal H -bundles $P_t \forall t \neq t_0$.

Following the procedure in Prop 6.6 we get a vector bundle $\mathcal{E}_{u|_{B_{\epsilon,u}}}/\mathcal{E}_{u|_{\partial B_{\epsilon,u}}}$ such that

$$c_2(\mathcal{E}_{u|_{B_{\epsilon,u}}}/\mathcal{E}_{u|_{\partial B_{\epsilon,u}}}) = \text{length}(\mathcal{E}_0^{**}/\mathcal{E}_0)_0$$

We now observe easily that the vector bundle $\mathcal{E}_{u|_{B_{\epsilon,u}}}/\mathcal{E}_{u|_{\partial B_{\epsilon,u}}}$ on S^4 also has a reduction of structure group to H . Thus we have a principal H -bundle P_ϵ on S^4 whose extension of structure group by $\rho : H \hookrightarrow SL(V)$ is $\mathcal{E}_{u|_{B_{\epsilon,u}}}/\mathcal{E}_{u|_{\partial B_{\epsilon,u}}}$.

By Prop 6.4 it follows that the second Chern class of $\mathcal{E}_{u|_{B_{\epsilon,u}}}/\mathcal{E}_{u|_{\partial B_{\epsilon,u}}}$ is a multiple of the Dynkin index m_ρ . Hence $\text{length}(\mathcal{E}_0^{**}/\mathcal{E}_0)_0$ is a multiple of m_ρ . This proves the theorem.

q.e.d

DEFINITION 6.8. Let $P_{t_0} \in \overline{M_H^0}$. Then we define the cycle $Z(P_{t_0})$ as:

$$Z(P_{t_0}) := \sum_{x \in X} \frac{\text{length}(\mathcal{E}_{t_0}^{**}/\mathcal{E}_{t_0})_x}{m_\rho} \cdot x$$

6.2. *Points of the moduli.* Let P be an H -quasibundle and let $P(\rho) = \mathcal{E}$. Further, let $\mathcal{E}_0$ be an associated graded sheaf of $\mathcal{E}$ and let us denote the double dual $\mathcal{E}_0^{**}$ by $\mathbb{E}$.

Let $Q(\mathbb{E}, l)$ be the quot scheme of torsion quotients of $\mathbb{E}$ of length l . By the choice of our $\mathbb{E}$, the torsion-free sheaves $F \in Q(\mathbb{E}, l)$ have all *reduction of structure groups* to quasibundles.

Notation: Let the induced total family quasibundles coming from $Q(\mathbb{E}, l)$ be denoted by $Q_H(\mathbb{E}, l)$. We thus have the following diagram:

$$\begin{array}{ccc} Q_H(\mathbb{E}, l) & \xrightarrow{f_\rho} & Q(\mathbb{E}, l) \\ \downarrow h_0 & \searrow h & \downarrow q \\ S^{l_0}(X) & \xrightarrow{''m_\rho''} & S^l(X) \end{array}$$

We note the following:

- (a) The degree l_0 is therefore $l_0 = \frac{l}{m_\rho}$.
- (b) The vertical map $h_0 : Q_H(\mathbb{E}, l) \rightarrow S^{l_0}(X)$ is the one which associates to each P the cycle $Z(P)$ defined above.
- (c) The map $''m_\rho'' : S^{l_0}(X) \rightarrow S^l(X)$ is the map induced by multiplication by the Dynkin index m_ρ which maps the cycle $Z(P)$ to $m_\rho \cdot Z(P) = Z(\mathcal{E}_0)$. Observe that this map is an *injection*.

Remark 6.9. (Action of $\text{Aut}(\mathbb{E})$ on the fibre of f_ρ)

Let $\alpha \in \text{Aut}(\mathbb{E})$. Let $\mathcal{E}$ be a μ -polystable torsion-free sheaf with $\mathbb{E} = \text{gr}^\mu(\mathcal{E})^{**}$. Let P be a quasibundle obtained from a reduction datum $\sigma : X \rightarrow S_{\mathcal{E}}//H$.

We first observe that since $\mathbb{E}$ is a locally free sheaf on X , by Hartogs theorem we see that the natural restriction map $\text{res}_U : \text{Aut}(\mathbb{E}) \rightarrow \text{Aut}(\mathbb{E}_U)$ is an isomorphism. (this holds when $\mathbb{E}$ is reflexive, cf. for example Cor 1.11.1 [26]).

By restricting α to U a big open set, we have an action of α on σ_U (restriction of σ to U). Let $\alpha \cdot \sigma_U = \sigma'_U$. Then by definition $\sigma'_U : U \rightarrow S_{\mathcal{E}}//H$ which extends *uniquely* to a new datum $\sigma' : X \rightarrow S_{\mathcal{E}}//H$. We define:

$$\alpha \cdot \sigma = \sigma'$$

It is clear that if P' is the quasibundle obtained from σ' , then $P'(\rho) = \mathcal{E}$ and $P^{**} \simeq P'^{**}$.

We then have the following:

PROPOSITION 6.10. *Let P and Q be two μ -polystable H -quasibundles in $Q_H(\mathbb{E}, l)$ lying in the same fibre of f_ρ . If $P^{**} \simeq Q^{**}$ then P and Q lie in the same connected component of the fibre of the map $h : Q_H(\mathbb{E}, l) \longrightarrow S^l(X)$ (or equivalently of the map $h_0 : Q_H(\mathbb{E}, l) \longrightarrow S^{l_0}(X)$).*

Proof: We first show that if P and Q are two *polystable* quasibundles in a fibre of the map $f_\rho : Q_H(\mathbb{E}, l) \longrightarrow Q(\mathbb{E}, l)$ and such that $P^{**} \simeq Q^{**}$ then P and Q lie in the same orbit of the action $Aut(\mathbb{E})$.

Note firstly that the polystability of P and Q as well as the property that they lie in the same fibre of f_ρ forces that the *associated torsion-free sheaves* in $Q(\mathbb{E}, l)$ are μ -polystable. Hence, both P and Q are *reductions* of structure groups of the “same” polystable torsion-free sheaf $\mathcal{E}_0$. One also sees that $\mathcal{E}_0^{**} \simeq \mathbb{E}$. In other words, $P, Q \in f_\rho^{-1}(\mathcal{E}_0)$.

Consider the restriction to a big open subset $U = U(\mathcal{E}_0)$. By definition, both P_U and Q_U are reductions of structure group of $\mathcal{E}_U \simeq \mathbb{E}_U$ and since $P^{**} \simeq Q^{**}$, these are isomorphic reductions of structure group of the principal bundle $\mathbb{E}_U$. Therefore they are in the same orbit of $Aut(\mathbb{E}_U)$. By the definition of the action in Remark 6.9, this implies that P and Q lie in the same orbit of $Aut(\mathbb{E})$.

Now since $\mathbb{E}$ is a polystable vector bundle on X (we confuse the principal bundle with the vector bundle associated to it) by an argument similar to the one in Lemma 5.15, it follows that for a general curve $C \in X$, $Aut(\mathbb{E}) \simeq Aut(\mathbb{E}|_C)$. Also by the Mehta-Ramanathan restriction theorem, $\mathbb{E}|_C$ is also a polystable vector bundle on C and hence its automorphism group is a product of $GL(n)$ ’s. Hence, $Aut(\mathbb{E})$ is connected. This implies that the orbit of $Aut(\mathbb{E})$ is connected and hence both P and Q lie in the same connected component of the fibre of h .

q.e.d

COROLLARY 6.11. *Let P and Q be two μ -semistable H -quasibundles in $Q_H(\mathbb{E}, l)$ lying in the same fibre of f_ρ . If $P_0^{**} \simeq Q_0^{**}$ then P and Q lie in the same connected component of the fibre of the map $h : Q_H(\mathbb{E}, l) \longrightarrow S^l(X)$.*

Proof: Observe that for any quasibundle P , an associated graded quasibundle is connected to P by a path (cf. Def 5.16). Hence the Corollary follows from the Prop 6.10.

PROPOSITION 6.12. *Let P and Q be two μ -polystable H -quasibundles in $Q_H(\mathbb{E}, l)$ lying in the same fibre of h . If further, $P^{**} \simeq Q^{**}$, then P and Q lie in the same connected component of the fibre of the map $h : Q_H(\mathbb{E}, l) \longrightarrow S^l(X)$ (or equivalently of the map $h_0 : Q_H(\mathbb{E}, l) \longrightarrow S^{l_0}(X)$).*

Proof: By the result of of Baranovsky ([5]) and Ellingsrud and Lehn ([14]) (and in rank 2, Li ([24, Prop 6.5])), one knows that the fibre of the map q , which we denote by T , is *connected*, in fact *irreducible*. In particular, every

torsion-free sheaf $\mathcal{F}$ in the fibre of q is a sheaf such that $\mathcal{F}^{**} \simeq \mathbb{E}$ and since $\mathbb{E}$ is polystable, it follows that so is $\mathcal{F}$. In other words, every closed point of the fibre T is represented by a polystable torsion-free sheaf.

Let us fix the polystable quasibundle P in the fibre of h and let Q be another polystable quasibundle such that $Q^{**} \simeq P^{**}$. The polystable torsion-free sheaves *associated* to P and Q , namely $\mathcal{E}$ and $\mathcal{F}$ are in T . If $\mathcal{E}$ and $\mathcal{F}$ coincide then the result follows from Prop 6.10. Let $f_\rho(P) = t_0$ and $f_\rho(Q) = t_1$. Since T is irreducible, without loss of generality we may take T to be an irreducible curve joining t_0 and t_1 . Let $\overline{T}$ be its normalisation. Hence, $\overline{T}$ is a smooth irreducible curve.

We claim that Q lies in the same connected component of the fibre of h which contains P .

Consider the family of torsion-free sheaves $\mathcal{E}_T$ (on $X \times T$) such that $\mathcal{E}_{t_0} \simeq \mathcal{E}$ and $\mathcal{E}_{t_1} \simeq \mathcal{F}$, which exists by the definition of the Quot scheme of torsion quotients. We shall work with the pull-back of $\mathcal{E}_T$ to $X \times \overline{T}$ and denote it by $\mathcal{E}_{\overline{T}}$. Thus we may take $t_0, t_1 \in \overline{T}$.

Since all the sheaves $\mathcal{E}_t$ lie over the same fibre T it follows that they are all polystable sheaves and have the same double dual $\mathbb{E}$. Let $\mathcal{U}$ be the maximal open subset where $\mathcal{E}_{\overline{T}}$ is locally free. Then we have an isomorphism of families on $\mathcal{U}$:

$$\mathcal{E}_{\overline{T}}|_{\mathcal{U}} \simeq \mathbb{E}_{\overline{T}}|_{\mathcal{U}}$$

where $\mathbb{E}_{\overline{T}} = \mathbb{E} \times \overline{T}$.

Since the quasibundle P is a “reduction of structure group” of $\mathcal{E}_{t_0}$, it implies that there is a section $\sigma : X \rightarrow (S_{\mathcal{E}_{t_0}})/H$.

Restricting this section to $\mathcal{U}_{t_0} = \mathcal{U} \cap (X \times t_0)$ we get a section $\sigma_{\mathcal{U}_{t_0}} : \mathcal{U}_{t_0} \rightarrow (S_{\mathcal{E}_{t_0}})/H$. Note that $\mathcal{U}_{t_0}$ is also a *big* open subset of X (cf. [35, 2.6, page 1187]).

Observe that the double dual principal bundle P^{**} of P is also obtained by a reduction of structure group of the principal bundle associated to $\mathbb{E}$.

Let us denote this reduction by $\tau : X \rightarrow (S_{\mathbb{E}})/H$. On the big open set $\mathcal{U}_{t_0}$ this reduction $\tau_{\mathcal{U}_{t_0}}$ and the reduction $\sigma_{\mathcal{U}_{t_0}}$ giving P are therefore mapped to each other by an automorphism $\phi \in \text{Aut}(\mathbb{E}|_{\mathcal{U}_{t_0}}) = \text{Aut}(\mathbb{E})$.

Since $\mathbb{E}_{\overline{T}} = \mathbb{E} \times \overline{T}$, the reduction section τ trivially extends to a section $\tau_{\overline{T}} : X \times \overline{T} \rightarrow (S_{\mathbb{E}_{\overline{T}}})/H$. Restricting $\tau_{\overline{T}}$ to $\mathcal{U}$ we get a section

$$\tau_{\mathcal{U}} : \mathcal{U} \rightarrow (S_{\mathbb{E}_{\overline{T}}|_{\mathcal{U}}})/H \simeq (S_{\mathcal{E}_{\overline{T}}})/H.$$

Since $\mathcal{U}$ is *big*, and since $X \times \overline{T}$ is *smooth (and hence normal)*, this section extends to give a new section $\overline{\tau} : X \times \overline{T} \rightarrow (S_{\mathcal{E}_{\overline{T}}|_{X \times \overline{T}}})/H$, i.e a “reduction datum” for the family $\mathcal{E}_{\overline{T}}$ (see Remark 2.3).

This gives a family $\overline{P}_{\overline{T}}$ of quasibundles on $X \times \overline{T}$ such that $\overline{P}_{t_0}$ and P are in the same orbit of $\text{Aut}(\mathbb{E})$ (by the element ϕ). Thus $\overline{P}_{t_0}$ and P lie in the same connected component of the fibre of h .

By the universal property of $Q_H(\mathbb{E}, l)$, we get a morphism induced by this family of quasibundles, namely

$$\psi : \overline{T} \longrightarrow Q_H(\mathbb{E}, l)$$

(ψ depends on the choice of the reduction τ).

Further, by the definition of the reduction datum on $\mathcal{U}$ all the quasibundles in the family $\overline{P}_{\overline{T}}$ (which are polystable) have *isomorphic double duals* P^{**} .

In particular, they lie in the same connected component of the fibre of h . Moreover, if Q corresponds to the point $t_1 \in \overline{T}$ then the quasibundles $\overline{P}_{t_1}$ and Q lie in the same fibre $f_\rho^{-1}(\mathcal{E}_{t_1})$. By Prop 6.10 they lie in the same orbit of $Aut(\mathbb{E})$ since $Q^{**} \simeq P^{**}$. This implies that the quasibundles P , $\overline{P}_{t_0}$, $\overline{P}_{t_1}$ and Q lie on the same connected component of the fibre of h . This proves the claim.

q.e.d

6.2.1. Characteristic classes of principal H -bundles. Let P be a principal H -bundle with H a simple algebraic group. Then by the tables in [8] for the “invariant degrees”, i.e the degrees of a generating set for the invariant polynomials under the adjoint action (or equivalently by the Chevalley restriction theorem, polynomials on the Cartan subalgebra invariant under the Weyl group action), we see that for H -simple, there is a *unique* generator of degree 2. Since we are over an algebraic surface, it follows that this invariant polynomial I_2 will be the only one which gives us a characteristic class in $H^4(X, \mathbb{Z})$. We shall denote this class by $c(P)$. (cf. [6])

Example 6.13. When H is classical, then $c(P) = c_2(P(V))$ where V is the defining representation.

We can now state the following key result:

PROPOSITION 6.14. *Let P and Q be two μ -semistable H -quasibundles. Let P_0^{**} be the canonical polystable principal bundle obtained from a choice of associated graded quasibundle of P . Let $c(P_0^{**}) = c(Q_0^{**})$ for the degree 2 characteristic class c of the principal bundles P_0^{**} and Q_0^{**} . Then P and Q define the same point of $M_H(\rho)$ if and only if we have an identification of pairs $(P_0^{**}, Z(P)) \simeq (Q_0^{**}, Z(Q))$ where $Z(P)$ is a cycle class in the symmetric power $S^{l_0}(X)$ given by*

$$Z(P) := \sum_{x \in X} \frac{\text{length}(\mathcal{E}_0^{**}/\mathcal{E}_0)_x}{m_\rho} \cdot x$$

with l_0 given by:

$$l_0 = \frac{c_2(\mathcal{E}_0) - c_2(\mathcal{E}_0^{**})}{m_\rho}$$

and where P (resp Q) is obtained from the torsion-free sheaf $\mathcal{E}$ (resp $\mathcal{F}$) by reduction of structure group via ρ and $\mathcal{E}_0$ (resp $\mathcal{F}_0$) is an associated graded sheaf of the semistable sheaf $\mathcal{E}$ (resp $\mathcal{F}$).

Proof: Let us assume that the quasibundles P and Q define the same point in the moduli space $M_H(\rho)$.

We first check that $P_0^{**} \simeq Q_0^{**}$: suppose that this does not hold. As we noted in Lemma 5.15, it implies that for a large degree curve C , the restrictions $P_0^{**}|_C \neq Q_0^{**}|_C$. By Lemma 5.5, we can separate the points corresponding to P and Q in $R_H(\rho)$ by invariant sections of $\mathcal{L}$ which contradicts the assumption that they define the same point in $M_H(\rho)$.

Again, the equality $Z(P) = Z(Q)$ follows immediately since the torsion-free sheaves $\mathcal{E}$ and $\mathcal{F}$ (from which the quasibundles P and Q are defined) give the same point in the moduli space M_G by ([19, Th 8.2.11]), and the cycle classes $m_\rho \cdot Z(P) = Z(\mathcal{E})$.

For the converse, suppose that we have an identification:

$$(P_0^{**}, Z(P)) \simeq (Q_0^{**}, Z(Q)).$$

By Cor 6.11, it is enough to check for polystable quasibundles in the fibre of h ; by Prop 6.12 it follows that P and Q lie in the *same connected component of the fibre of $h : Q_H(\mathbb{E}, l) \rightarrow S^l(X)$* .

By the definition of the determinant line bundle on $R_H(\rho)$, it is obtained by pulling back the corresponding line bundle from the total family R_G . By the theorem of Baranovsky ([5]) and Ellingsrud and Lehn ([14]) one knows that the fibres of the morphism $q : Q(\mathbb{E}, l) \rightarrow S^l(X)$ are *connected* and also that the line bundle $\mathcal{L}$ restricted to this fibre is *trivial* (cf. [19, 8.2.1] and Li ([24, Lemma 3.4, 3.5])).

By the commutative diagram seen earlier, we have $q \circ f_\rho = h$. Therefore $\mathcal{L}$ is trivial on the fibres of h as well. In particular, the connected components of the fibre of h get mapped to the same point in $M_H(\rho)$. This proves that the points defined by P and Q coincide in $M_H(\rho)$.

q.e.d

PROPOSITION 6.15. *There is a canonical morphism $j : M_H(\mathfrak{c})^s \rightarrow M_H(\rho)$ which is an embedding of the moduli space of isomorphism classes μ -stable principal H -bundles in $M_H(\rho)$.*

Proof: The existence of the morphism j follows by the weak coarse moduli property of the moduli space $M_H(\rho)$. The injectivity of j follows from Lemma 5.15.

Now choose a general curve $C \in |a\Theta|$ which gives Bogomolov's effective restriction theorem, it is easily seen that the differential of the restriction map $r|_C$ (by usual arguments involving Enriques-Severi lemma) is *injective*. This immediately implies that the map j is an *embedding*.

q.e.d

Let $\mathfrak{M}_H$ be defined by the disjoint union:

$$\mathfrak{M}_H = \coprod_{l \geq 0} M_H^{\mu\text{-poly}}(\mathfrak{c} - l) \times S^l(X) \quad (*)$$

where $M_H^{\mu-pol\gamma}(\mathbf{c} - l)$ is the moduli space of μ -semistable principal H -bundles with characteristic class $\mathbf{c} - l$ (represented as classes of polystable bundles); in particular, the big stratum is $M_H^{\mu-pol\gamma}(\mathbf{c}) = M_H^0$.

In conclusion we have the following:

THEOREM 6.16. *There is a set-theoretic inclusion*

$$\mathfrak{M}_H \subset M_H(\rho)(\mathbb{C})$$

Furthermore, the underlying set of points $\overline{M_H^0}(\mathbb{C})$, of the closure of the moduli space of equivalence classes of μ -semistable principal H -bundles, is a subset of $\mathfrak{M}_H$. In fact, $\overline{M_H^0}(\mathbb{C})$ is independent of the representation ρ upto homeomorphism.

Proof: To complete the proof of the theorem we need only show that if (P^{**}, Z) is a point on the right hand side of (*) above, then there exists a point in $M_H(\rho)$ which corresponds to it.

Since P^{**} is a polystable principal H -bundle, using ρ if we extend the structure group to $G = GL(V)$, we get a μ -polystable locally free sheaf $\mathbb{E} \in M_G$ together with a reduction $\sigma : X \rightarrow \mathbb{E}(G/H)$ (as before we confuse the locally free sheaf $\mathbb{E}$ with the associated principal $GL(V)$ -bundle $\mathbb{E}$).

By the results of Li, it is known that the map $j : S^l(X) \rightarrow M_G$, induced by the morphism from $Q(\mathbb{E}, l) \rightarrow M_G$, is an *embedding*. Therefore the element $m_\rho \cdot Z \in S^l(X)$ gives a polystable torsion-free sheaf $\mathcal{E}$ such that $\mathcal{E}^{**} \simeq \mathbb{E}$ and such that $m_\rho \cdot Z = Z(\mathcal{E})$.

We therefore have a reduction of structure group of $\mathcal{E}|_{U(\mathcal{E})}$ coming from the restriction $\sigma|_{U(\mathcal{E})}$ of the reduction of structure group of $\mathbb{E}$ which gives the principal bundle P^{**} . Viewing this $\sigma|_{U(\mathcal{E})}$ as a morphism $U(\mathcal{E}) \rightarrow S_{\mathcal{E}}//H$, we get a canonical extension $\sigma_1 : X \rightarrow S_{\mathcal{E}}//H$. This gives rise to a polystable H -quasibundle $P \in M_H(\rho)$ which corresponds to the point (P^{**}, Z) . That this map which sends $(P^{**}, Z) \rightarrow P$ is injective follows from the first part of the proof of Prop 6.14.

The last part of the theorem which realises the closure $\overline{M_H^0}(\mathbb{C})$ in $\mathfrak{M}_H$ follows from the consequences of Theorem 6.7 and the definition Def 6.8. The uniqueness upto homeomorphism is precisely the content of Prop 6.17 below.

q.e.d

PROPOSITION 6.17. *The closure $\overline{M_H^0}(\mathbb{C})$ is unique upto homeomorphism. In fact, the isomorphic copies of M_H^0 defines a correspondence between the closures which gives the homeomorphism. Furthermore, the normalisations of the closures are isomorphic as reduced projective schemes.*

Proof: Since most the points needed in the proof have already been discussed above, we content ourselves by giving the main steps in the proof of this Proposition.

Let $\rho_i : H \hookrightarrow SL(V_i)$ for $i = 1, 2$ be two faithful representations of H and suppose that $P_t(i)$ be family of μ -semistable principal H -bundles with isomorphic generic fibres and with two limits as *polystable H -quasibundles* coming from the representations ρ_i . Let the limits be $P(i)$, $i = 1, 2$.

Claim 1: The double dual polystable H -bundles $P(1)^{**} \simeq P(2)^{**}$. To see this, restrict the families $P_t(i)$ to a general high degree curve C which avoids the singular loci of $P(i)$ for $i = 1, 2$. Then, since for such families over curves the (polystable) limit is uniquely defined, it follows that $P(1)^{**}|_C \simeq P(2)^{**}|_C$. By Lemma 5.15 the claim follows.

Claim 2: The cycles $Z(P(i))$ also coincide. This follows from two observations, namely that for the limits the open subsets where the $P(i)$ are genuine principal bundles coincide and hence the sets $Sing(P(i))_{red}$ coincide. The multiplicities at all the singular points also coincide by the discussion before Definition 6.8.

These two claims imply the set-theoretic identification of the closures. Moreover, the above discussion gives an identification of closures of any curve in M_H^0 . This implies that the projections to $M_H(\rho_i)$ from the graph $\Gamma \subset M_H(\rho_1) \times M_H(\rho_2)$, which is closed in the product, actually give homeomorphisms. The comment on normalisations now follows by Zariski's main theorem.

COROLLARY 6.18. *The closure of subset $M_H(c)^s \subset M_H^0$ gives the Donaldson-Uhlenbeck compactification of the moduli space of principal H -bundles with irreducible ASD connections.*

6.2.2. Relationship with Gómez-Sols and Schmitt's moduli space. We work in the set-up of Schmitt's recent paper (cf. [35] and [36]). Fix a faithful representation $\rho : H \hookrightarrow SL(V)$. Define the moduli functors

$$\underline{M}(\rho)_P^{ss} : \underline{Sch}_{\mathbb{C}} \longrightarrow \underline{Set}$$

which sends S to “equivalence classes of families of Gieseker–Maruyama semistable honest singular principal H -bundles (or H -quasibundles) with Hilbert polynomial P on X parametrised by S ”. Then the main theorem of [36] is that there is a projective scheme $\mathcal{M}(\rho)_P^{ss}$ which *coarsely represents* the functor $\underline{M}(\rho)_P^{ss}$. The notions of Gieseker–Maruyama semistability of quasibundles is defined in [18] and [36] and it is also shown that if $P = (\mathcal{E}, \tau)$ is an H -quasibundle then the following equivalence holds ([36, Section 5.1]): P is μ -stable $\implies P$ is Gieseker – Maruyama – stable $\implies P$ is Gieseker – Maruyama – semistable $\implies P$ is μ -semistable. These implications, together with the coarse moduli property of the functor $\underline{M}(\rho)_P^{ss}$ implies that there is a morphism:

$$\mathcal{M}(\rho)_P^{ss} \longrightarrow M_H(\rho).$$

At the risk of repetition, we now summarise the above results in the following:

THEOREM 6.19. *Let H be a semisimple algebraic group.*

- (a) *There exists a projective scheme $M_H(\rho)$ which parametrises equivalence classes of μ -semistable H -quasibundles with fixed characteristic classes, on the smooth projective surface X .*
- (b) *This has an open subscheme of equivalence classes μ -semistable principal H -bundles M_H^0 .*
- (c) *If M_H^s is the subscheme of $M_H(\rho)$ consisting of the stable principal H -bundles, then its closure $\overline{M^s}$ in $M_H(\rho)$ corresponds to the Donaldson-Uhlenbeck compactification of the moduli space of μ -stable vector bundles on X (or equivalently of anti self-dual connections).*
- (d) *When H is a simple algebraic group the points of the moduli space $\overline{M_H^0}$ are given by the pairs $(P_0^{**}, Z(P))$ where, to each semistable H -quasi bundle P , we associate P_0^{**} the canonical polystable principal H -bundle obtained from the quasibundle P and the cycle class $Z(P)$ which lies in $S^{l_0}(X)$, l_0 being given by:*

$$\frac{c_2(\mathcal{E}_0) - c_2(\mathcal{E}_0^{**})}{m_\rho}.$$

Note that P is obtained from the torsion-free sheaf $\mathcal{E}$ by reduction of structure group via ρ and $\mathcal{E}_0 = gr^\mu(\mathcal{E})$ and m_ρ is the Dynkin index associated to ρ .

- (e) *When H is a simple algebraic group the underlying set of points of the moduli space $\overline{M_H^0}$, upto homeomorphism, is independent of the representation ρ .*
- (f) *By ([36, Main theorem]), we have a morphism*

$$\mathcal{M}(\rho)_P^{ss} \longrightarrow M_H(\rho)$$

- (g) *In a certain sense, $\overline{M_H^0}$ is the minimal compactification of M_H^0 .*

7. NON-EMPTYNESS OF THE MODULI SPACE

In this section H is an arbitrary semisimple algebraic group. The aim of this section is to prove that the moduli space $M_H(c)^s$ of μ -stable principal H -bundles on a smooth projective surface X for *large* characteristic classes c is non-empty. When $H = SL(2)$ this is a theorem due to Taubes [40] and later due to Gieseker [16]. We quote their result:

THEOREM 7.1. *(Taubes, Gieseker) For any choice of polarisation Θ , for all $c \geq 2p_g(X) + 2$, the moduli space $M_{SL(2)}(c)^s$ of stable rank 2 bundles with trivial determinant and $c_2 = c$ is non-empty.*

7.1. Stable bundles and the principal three dimensional subgroup. We begin by recalling a few facts about semisimple Lie algebras. Let $\mathfrak{h} = Lie(H)$. Then one knows that there exist the so-called *principal* $\mathfrak{sl}(2)$'s inside $\mathfrak{h}$ which are distinguished by the property that they do not lie in any parabolic

subalgebra of $\mathfrak{h}$. This gives a representation $\phi : SL(2) \rightarrow H$ which is *irreducible*, in the sense that $Im(\phi)$ is not contained in any proper parabolic subgroup of H . (cf. [8] Chapter 8, Exercise 5 §11, page 246).

We fix one such homomorphism of a principal $SL(2) \rightarrow H$.

Let C be a smooth projective curve of genus $g \geq 2$. Let V be a stable vector bundle of rank 2. Then by the Narasimhan–Seshadri theorem one knows that there exists an irreducible representation $\rho : \pi_1(C) \rightarrow SU(2)$ such that $V \simeq V_\rho$.

DEFINITION 7.2. We define the monodromy subgroup $\mathcal{M}(V)$ of $V \simeq V_\rho$ to be the Zariski closure of $Im(\rho)$ in $SL(2)$. It can be viewed as the minimal subgroup to which the structure group of V can be reduced.

Remark 7.3. Recall that one knows that the *reductivity* of $\mathcal{M}(V)$ is equivalent to the polystability of V . Since one knows the set of all finite subgroups of $SL(2)$ (this is classical, for example cf. [38]), the only possibilities for $\mathcal{M}(V)$ are the following:

- (a) Finite cyclic groups C_n and the dihedral groups D_n .
- (b) The alternating groups A_4 and A_5 and the permutation group S_5 .
- (c) The whole of $SL(2)$ or its maximal torus.

Of these, since V is stable, we can omit the maximal torus and the cyclic groups C_n . So $\mathcal{M}(V)$ can either be the alternating groups, S_5 , or the dihedral groups apart from the whole of $SL(2)$.

We wish to estimate the set Z_C of representations of $\pi_1(C)$ in $SL(2)$ which lie entirely in these families of finite groups upto conjugacy by the diagonal action of $SL(2)$.

It is not hard to see that since $\pi_1(C)$ is given by $2g$ generators with a single relation, this set Z is at most *countable*. This implies the following lemma.

LEMMA 7.4. *The locus of points in the moduli space of stable vector bundles $M_{C,SL(2)}$ of rank 2 and trivial determinant on the curve C whose monodromy is among the set of finite subgroups listed above has countable cardinality.*

Using this we have the following:

PROPOSITION 7.5. *There exists a rank 2 stable bundle E with $c_2(E) \gg 0$ and trivial determinant on the surface X such that the restriction $E|_C$ to general curve $C \subset X$ of high degree has monodromy subgroup to be the whole of $SL(2)$ itself.*

Proof: For $c = c_2(E) \gg 0$ (by Theorem 7.1), it is known that the moduli space $M_X(2, \mathcal{O}, c_2)^s = M_{SL(2)}(c)^s$ is non-empty and has dimension $d(c) = 4c - 3\chi(\mathcal{O}_X)$. Since $c_2(E) \gg 0$, it follows that we can make $d \geq 1$.

Now we consider the restriction map $r_C : M_{SL(2)}(c)^s \rightarrow M_{C,SL(2)}^s$. If the curve C is chosen sufficiently general and high degree k then by the Mehta–Ramanathan theorem and [15, Prop 2.2], it can be seen that for $k \gg 0$,

the map r_C is in fact an *immersion*. This implies that $Im(r_C) \subset M_{C,SL(2)}^s$ is not completely contained in the subset Z_C of stable bundles with finite monodromy groups. Therefore there exists at least one stable bundle with the whole of $SL(2)$ as its monodromy subgroup. This proves the proposition.

Before we proceed to our next lemma we recall the following formulation of *irreducible* sets ([31, page 129]):

DEFINITION 7.6. Let K be a maximal compact subgroup of H . A subset $A \subset K$ is said to be *irreducible* if

$$\{Y \in \mathfrak{h} | ad x(Y) = Y, \forall x \in A\} = \text{centre of } \mathfrak{h}.$$

A (homomorphism) representation $\delta : \Gamma \longrightarrow K$ is said to be irreducible if the image $\delta(\Gamma)$ is irreducible.

Note that in our case since H is semisimple *centre of* $\mathfrak{h}$ is trivial. We have the following equivalences and we quote:

PROPOSITION 7.7. ([31, Prop 2.1]) Let K be a maximal compact subgroup of H and let $A \subset K$ be a subgroup. Then the following are equivalent

- (a) A is irreducible.
- (b) A leaves no parabolic subalgebra of $\mathfrak{h}$ invariant.
- (c) For any parabolic subgroup $P \subset H$, A acts without fixed points on H/P .

LEMMA 7.8. If V is a stable bundle on a curve C such that $\mathcal{M}(V) = SL(2)$ then $V(\phi)$ is a stable H -bundle on C by an extension of structure group via the principal homomorphism ϕ .

Proof: Let $\rho : \pi_C \longrightarrow SU(2)$ is an irreducible unitary representation and let $V \simeq V_\rho$ be a stable bundle on C such that $\mathcal{M}(V) = SL(2)$.

This implies that $Im(\rho)$ is Zariski dense in $SL(2)$. Since $Im(\phi)$ is not completely contained in any parabolic subgroup of H by Prop 7.7 we have the following:

$$\{Y \in \mathfrak{h} | ad x(Y) = Y, \forall x \in \mathfrak{sl}(2)\} = \text{centre of } \mathfrak{h} = \text{trivial}.$$

In other words, we can say that $Im(\phi)$ is an *irreducible subset* in H (or equivalently, by the “unitarian trick”, $Im(\phi|_{\mathfrak{su}(2)})$ is an *irreducible* subset of K , where as above, $K \subset H$ is a maximal compact subgroup).

By [31, Prop 2.2], to show that $V(\phi)$ is stable as an H -bundle, we need only show that the representation

$$\eta = (\phi \circ \rho) : \pi_1(C) \longrightarrow K$$

is *irreducible*.

If $Y \in \mathfrak{h}$ is such that $ad x(Y) = Y, \forall x \in Lie(Im(\eta))$, then by the density of $Im(\eta)$ in $Im(\phi) = \mathfrak{sl}(2)$ and hence by continuity, we see that for such an element Y , one has $adx(Y) = Y, \forall x \in \mathfrak{sl}(2)$ and hence $Y \in \text{centre of } \mathfrak{h}$. This shows that $Im(\eta)$ is an irreducible set in $K \subset H$ and we are done. *q.e.d*

We can now conclude the following:

THEOREM 7.9. *The moduli space $M_H(c)^s$ of μ -stable principal H -bundles on a smooth projective surface X is non-empty for $c > \delta$, where δ depends only of $p_g(X)$ and not on the polarisation Θ on X .*

Proof: We claim that if $\phi : SL(2) \longrightarrow H$ is as above a principal $SL(2)$ in H then any E as in Prop 7.5 has the property that $E(\phi)$ is a *stable* principal H -bundle.

By the converse to the Mehta–Ramanathan theorem (Lemma 4.5), we see that it is enough to prove that $E(\phi)|_C$ is *stable*. Since $\mathcal{M}(E|_C) = SL(2)$ this is immediate by Lemma 7.8. The *largeness* of the characteristic classes of the associated H -bundle is determined by the largeness of the c_2 of the rank 2 bundle E . This can be determined by the general methods of Borel and Hirzebruch (see the recent preprint Beauville [6] for this).

q.e.d

7.2. Concluding remarks.

- (a) The questions addressed in this paper can be posed in positive characteristic as well, but as in the case of curves a subtler analysis of representation theoretic bounds such as the ones considered in [4] will have to be carried out. We hope to do this in future.
- (b) The basic questions regarding irreducibility, reducedness, generic smoothness and normality are yet to be answered for these moduli spaces.

REFERENCES

- [1] M.F. Atiyah and R. Bott: The Yang-Mills equations over Riemann Surfaces, *Phil. Tr. R. Soc. Lond. A* **308** (1982), 523-615.
- [2] M. Atiyah, N. Hitchin and I.M. Singer : Self-duality in four dimensions, *Proc. Royal Soc. Lond A* **362** (1978), 425-461.
- [3] V. Balaji and C.S. Seshadri: Semistable Principal Bundles-I (in characteristic zero), *Journal of Algebra* **258** (2002), 321-347.
- [4] V. Balaji and A.J. Parameswaran : Semistable principal bundles-II (in positive characteristics). *Transformation Groups*, **18**, No 1, (2003), pp 3-36.
- [5] V. Baranovsky : Moduli of sheaves on surfaces and action of the oscillator algebra, *J. Diff. Geom.*, **55**, No 2, (2000), pp 193-227.
- [6] A. Beauville : Chern classes for representations of reductive groups, *math.AG/0104031*.
- [7] I. Biswas and T.L. Gómez : Restriction theorem for principal bundles, *Math. Annalen*, **327**, (2003), 773-792.
- [8] N. Bourbaki : Groupes et algèbres de Lie, Chap 7-8, Paris, Hermann, (1975).
- [9] A. Braverman, M. Finkelberg and D. Gaitsgory : Uhlenbeck spaces via affine Lie algebras, *math.AG/0301176*.
- [10] J-L Colliot-Thélène and J. Sansuc : Fibrés quadratiques et composantes connexes réelles, *Math. Annalen*, **244** (1979) pp 105-134.
- [11] S.K. Donaldson : Polynomial invariants for smooth four manifolds, *Topology* **29**, (1990), 257-315.
- [12] S.K. Donaldson and P.B. Kronheimer: The Geometry of Four-Manifolds, Oxford (1990).

- [13] E.B.Dynkin : Semisimple subalgebras of semisimple Lie algebras, *Am Math Soc Transl Ser II* **6**, (1957) 111-244.
- [14] G.Ellingsrud and M.Lehn : Irreducibility of the punctual quotient scheme of a surface. *Ark. Mat.* **37** (1999), no. 2, 245–254.
- [15] R.Friedman and J.Morgan : Smooth four-manifolds and complex surfaces, *Ergebnisse*, Vol 27, Springer Verlag (1994).
- [16] D.Gieseker : A construction of stable bundles on an algebraic surface, *J. Diff. Geom* **27**, (1988), 137-154.
- [17] D.Gieseker and J.Li : Moduli of high rank vector bundles over surfaces, *J.AMS* **9**, (1996), 107-151.
- [18] T.Gómez and I.Sols : Moduli spaces of principal sheaves over projective varieties, *Annals of Math.* (to appear).
- [19] D.Huybrechts and M.Lehn: The Geometry of moduli spaces of sheaves, *Aspects of Mathematics E31*, Vieweg, Braunschweig/Wiesbaden, (1997).
- [20] A.Langer: Semistable sheaves in positive characteristic, *Annals of Math.*
- [21] S.Kumar, M.S.Narasimhan and A.Ramanathan, Infinite Grassmanians and moduli spaces of G -bundles, *Math Ann* **300**(1994), 41-75.
- [22] S.Langton: Valutive criterion for families of vector bundles on algebraic varieties, *Annals of Mathematics*(2) **101** (1975) pp 88-110.
- [23] J.Le Potier: Fibré déterminant et courbes de saut sur les surfaces algébriques, *Complex projective Geometry*, London Mathematical Society : Bergen(1989), 213-240.
- [24] J.Li: Algebraic geometric interpretation of Donaldson's polynomial invariants of algebraic surfaces, *J.Diff.Geom* **37** (1993), 416-466.
- [25] J.Li: Compactification of moduli of vector bundles over algebraic surfaces, (In a volume of papers dedicated to Prof C.G.Gu).
- [26] M.Maruyama: The Theorem of Grauert-Mülich-Spindler, *Math. Ann* **255**, (1981) 317–333.
- [27] V.B.Mehta and A.Ramanathan: Restriction of stable sheaves and representations of the fundamental group, *Invent.Math* **77**, (1984), 163–172.
- [28] J.Morgan : Comparison of the Donaldson polynomial invariants with their algebro geometric analogues, *Topology* **32** (1993), 449-488.
- [29] M.S. Narasimhan and C.S.Seshadri: Stable and unitary vector bundles on a compact Riemann surface, *Annals of Mathematics* (2)**82** (1965) pp 540-567.
- [30] P.Newstead: Introduction to moduli problems, TIFR Lecture Notes in Mathematics, (1978).
- [31] A.Ramanathan: Stable principal bundles on a compact Riemann surface, *Math. Ann*, **213**, (1975) 129-152.
- [32] A. Ramanathan: Stable principal bundles on a compact Riemann surface - Construction of moduli space (*Thesis*, Bombay University 1976) *Proc.Ind.Acad.Sci* **106**, (1996), 301-328 and 421-449.
- [33] A.Ramanathan and S.Subramaniam : Einstein–Hermitian connections on principal bundles and stability, *J.Reine Angew. Math* **390** (1988), 21-31.
- [34] S.Ramanan and A.Ramanathan : Some remarks on the instability flag, *Tohoku Math. Journ* **36** (1984), 269-291.
- [35] A.Schmitt: Singular Principal Bundles over higher dimensional manifolds and their moduli spaces, *IMRN*, **23** (2002), 1183-1209.
- [36] A.Schmitt: A closer look at semistability for singular principal bundles, (to appear in *IMRN*).
- [37] J.P.Serre: Espaces fibrés algébriques, Anneaux de Chow et applications, Séminaire Chevalley, (1958), also *Documents Mathématiques* **1** (2001), 107-140.
- [38] J.P.Serre: Moursund Lectures, University of Oregon Mathematics Department (1998).

- [39] C. Simpson: Higgs bundles and Local systems, *Pub. I.H.E.S.* **75** (1992), 5-95.
- [40] C.H.Taubes : Self-dual connections on 4-manifolds with indefinite intersection matrix, *J.Diff.Geom* **19** (1984), 517-560.

V.BALAJI CHENNAI MATHEMATICAL INSTITUTE 92, G.N.CHETTY ROAD, CHENNAI-600017 INDIA

E-mail address: `balaji@cmi.ac.in`